

## Announcements and Such

- Today's Music: *The Doors*
- I have posted my solutions to HW #4 and HW #5.
- HW #6 is due on Thursday @ 4pm.
- ☞ **The final is *in class* on Thursday. You'll be given 3 hours to do it.**
- I've posted two important handouts concerning the final exam:
  - The (Complete) Natural Deduction Rules Handout (provided at final).
  - A sample final exam, which has the same structure as the actual final. This sample was discussed, in detail, in lecture last week.
- Today: Chapters 7 & 8 — L2PL
  - I will only be covering (some of) the **L2PL** parts of Chapters 7 & 8.
  - I will say something about what will be on the final at the end of today's lecture (which, alas, will be my last lecture at Berkeley).

## Why (†) is So Important — L2PL vs LMPL: Infinite Domains

- In LMPL, if  $p$  is true on any interpretation  $\mathcal{I}$ , then it is true on a *finite* interpretation. Indeed,  $p$  will be true on an interpretation of size no greater than  $2^k$ , where  $k$  is the # of monadic predicate letters in  $p$ .
- In L2PL, some statements are true *only* on *infinite* interpretations. It is for this reason that there is no general decision procedure for validity (or logical truth) in L2PL. (†) on the last slide is a good example of this.

$$(†) \quad (\forall x)(\exists y)Rxy, (\forall x)(\forall y)(\forall z)[(Rxy \ \& \ Ryz) \rightarrow Rxz] \neq (\exists x)Rxx$$


- **Fact.** *Only infinite interpretations  $\mathcal{I}$  can be counterexamples to the validity in (†).* To see why, try to *construct* such an interpretation.
- We start by showing that no interpretation  $\mathcal{I}_1$  with a 1-element domain can be an interpretation on which the premises of (†) are  $\top$  and its conclusion is  $\perp$ . Then, we will repeat this argument for  $\mathcal{I}_2$  and  $\mathcal{I}_3$ .
- This reasoning can, in fact, be shown correct for *all* (finite)  $n$ . So, only  $\mathcal{I}$ 's with infinite domains will work [*e.g.*,  $\mathcal{D} = \mathbb{N}$ ,  $Rxy: x < y$ ].
- Begin with a 1-element domain  $\{\alpha\}$ . For the conclusion of (4) to be  $\perp$ , no

object can be related to itself:  $(\forall x) \sim Rxx$ . Thus, we must have  $\sim Raa$ :



$\alpha$


- But, to make the first premise  $\top$ , we need there to be *some*  $y$  such that  $Ray$  is  $\top$ . That means we need *another object*  $\beta$  to allow  $Rab$ . Thus:



$\alpha$

$\beta$


- Now, because we need the conclusion to remain  $\perp$ , we must have  $\sim Rbb$ . And, because we need the first premise to remain  $\top$ , we need there to be *some*  $y$  such that  $Rby$  is  $\top$ . We could *try* to make  $Rba$   $\top$ , as follows:




$\alpha$

$\beta$

- But, this picture is not consistent with the second premise being  $\top$  and (at the same time) the conclusion being  $\perp$ . If  $R$  is transitive, then  $Rab \ \& \ Rba$  (as pictured) entails  $Raa$ , which makes the conclusion  $\top$ .




- Thus, the only way to consistently ensure that there is some  $y$  such that  $Rby$  is to introduce *yet another object*  $y$  (such that  $Rbc$ ), which yields:



- Again, in order to make the conclusion  $\perp$ , we must have  $\sim Rcc$ , and in order to make the first premise  $\top$ , there must be some  $y$  such that  $Rcy$ .
- We could *try* to make either  $Rca$  or  $Rcb$  true. But, both of these choices will end-up with the same sort of inconsistency we just saw with  $\beta$ .

- In other words, *no finite interpretation* will give us what we want here.
- However, if we let  $\mathcal{D} = \mathbb{N}$  and  $Rxy: x < y$ , then we get what we want.



- That is, the relation  $Rxy: x < y$  on the natural numbers  $\mathbb{N}$  is such that:
  - For all  $x$ , there exists a  $y$  such that  $x < y$ . [seriality]
  - For all  $x, y, z$ , if  $x < y$  and  $y < z$ , then  $x < z$ . [transitivity]
  - For all  $x$ ,  $x \not< x$ . [irreflexivity]
- It is crucial that the set  $\mathbb{N}$  of *all* natural numbers is *infinite*. The relation  $<$  cannot satisfy all three of these properties on *any finite* domain.
- *I.e.*, no finite subset of  $\mathbb{N}$  will suffice to show that the invalidity in (4) holds. Equivalently, the following sentence of L2PL is  $\perp$  on *all finite*  $\mathcal{I}$ 's:
 
$$p \stackrel{\text{def}}{=} (\forall x)(\exists y)Rxy \ \& \ (\forall x)(\forall y)(\forall z)[(Rxy \ \& \ Ryz) \rightarrow Rxz] \ \& \ (\forall x) \sim Rxx$$
- This sort of thing *cannot happen* in LMPL. In this sense, the introduction of a single 2-place predicate involves a *quantum leap* in complexity.

## Some Further Remarks on Validity in L2PL

- As I just explained, there is no general decision procedure for  $\models$  claims in L2PL. This is because we can't always establish  $\neq$  claims in finite time.
- However, there is a method for proving  $\models$  claims — *natural deduction*. And, L2PL's natural deduction system *is exactly the same as LMPL's!*
- Before we get to proofs, however, I want to look at the alternating quantifier example that I said separates LMPL and L2PL.
- As we have seen,  $(\forall x)(\exists y)Rxy \neq (\exists y)(\forall x)Rxy$ . But, the converse entailment *does* hold. That is,  $(\exists y)(\forall x)Rxy \models (\forall x)(\exists y)Rxy$ .
- We will *prove* — *i.e., deduce* —  $(\exists y)(\forall x)Rxy \vdash (\forall x)(\exists y)Rxy$  shortly.
- Before we do that, let's think about  $(\exists y)(\forall x)Rxy \models (\forall x)(\exists y)Rxy$  using our definitions, and our informal method of thinking of  $\forall$  as & and  $\exists$  as  $\vee$ . This is interesting for both directions of the entailment.
- But, we need to be much more careful here than with LMPL!

- First, consider what  $(\exists y)(\forall x)Rxy$  says on a domain of size  $n$ :  

$$(\exists y)(\forall x)Rxy \approx_n (\forall x)Rxa \vee (\forall x)Rxb \vee \dots \vee (\forall x)Rxn$$

$$\approx_n (Raa \& \dots \& Rna) \vee (Rab \& \dots \& Rnb) \vee \dots \vee (Ran \& \dots \& Rnn)$$
- Next, consider what  $(\forall x)(\exists y)Rxy$  says on a domain of size  $n$ :  

$$(\forall x)(\exists y)Rxy \approx_n (\exists y)Ray \& (\exists y)Rby \& \dots \& (\exists y)Rny$$

$$\approx_n (Raa \vee \dots \vee Ran) \& (Rba \vee \dots \vee Rbn) \& \dots \& (Rna \vee \dots \vee Rnn)$$
- Then, we notice that these two sentential forms are intimately related. Specifically, we note that  $(\exists y)(\forall x)Rxy$  has the following  $n$ -form:  

$$\mathcal{X}_n = (p_1 \& p_2 \& \dots \& p_n) \vee (q_1 \& q_2 \& \dots \& q_n) \vee \dots \vee (r_1 \& r_2 \& \dots \& r_n)$$
- And, we notice that  $(\forall x)(\exists y)Rxy$  has the following  $n$ -form:  

$$\mathcal{Y}_n = (p_1 \vee q_1 \vee \dots \vee r_1) \& (p_2 \vee q_2 \vee \dots \vee r_2) \& \dots \& (p_n \vee q_n \vee \dots \vee r_n)$$
- **Fact.**  $\mathcal{X}_n \models \mathcal{Y}_n$ , for any  $n$ . Each disjunct of  $\mathcal{X}_n$  entails every conjunct of  $\mathcal{Y}_n$ . **Caution!** This *doesn't* show that  $(\exists y)(\forall x)Rxy \models (\forall x)(\exists y)Rxy$ !
- **Fact.**  $\mathcal{Y}_n \not\models \mathcal{X}_n$ , for all  $n > 1$ . This can be shown (next slide) using only LSL reasoning. This *does* show that  $(\forall x)(\exists y)Rxy \not\models (\exists y)(\forall x)Rxy$ .
- The moral is that our “informal” semantical approach to the quantifiers works for LMPL, since no infinite domains are required for  $\neq$  in LMPL.

- However, our “informal” semantical approach breaks down for L2PL, since we sometimes need an infinite domain to establish  $\neq$  in L2PL.
- In L2PL, if the “informal” method above reveals  $p_n \neq q_n$  for *some* finite  $n$ , then it *does* follow that  $p \neq q$ . For instance,  $\mathcal{Y}_2 \neq \mathcal{X}_2$  on the last slide:
  - $(Raa \vee Rab) \& (Rba \vee Rbb) \neq (Raa \& Rba) \vee (Rab \& Rbb)$
  - This is just an LSL problem with 4-atoms [ $A = Raa$ ,  $B = Rab$ ,  $C = Rba$ ,  $D = Rbb$ ]. Truth-tables will generate a counterexample.
- On the other hand, if (in L2PL) our “informal” method indicates (as above) that  $p_n \models q_n$  for *all* finite  $n$ , this does *not* guarantee  $p \models q$ . *E.g.*:
  - $p = (\forall x)(\exists y)Rxy \& (\forall x)(\forall y)(\forall z)[(Rxy \& Ryz) \rightarrow Rxz]$ .
  - $q = (\exists x)Rxx$ .
- We showed above (informally) that  $p_n \models q_n$  for *all* finite  $n$ . But, we also saw that there are infinite interpretations on which  $p$  is  $\top$  but  $q$  is  $\perp$ .

## Natural Deduction in L2PL

- The ND system we already have is sound and complete for L2PL (140A!).
- So, we're already in a position to do natural deductions in L2PL.
- Here are four examples, with solutions on subsequent slides:
  1.  $(\exists y)(\forall x)Rxy \vdash (\forall x)(\exists y)Rxy$ 
    - This is the alternating quantifier example we just examined.
  2.  $(\forall x)(\forall y)(\forall z)[(Rxz \& Ryz) \rightarrow Rxy], (\forall x)Rxx \vdash (\forall x)(\forall y)(Rxy \rightarrow Ryx)$ 
    - This is: Euclidean-ness + reflexivity entails symmetry.
  3.  $(\forall x)(\forall y)(\forall z)[(Rxz \& Ryz) \rightarrow Rxy], (\forall x)Rxx$   
 $\vdash (\forall x)(\forall y)(Rxy \rightarrow Ryx)$ 
    - This is: Euclidean-ness + reflexivity entails transitivity.
  4.  $(\forall x)Fx \leftrightarrow \sim(\exists x)(\exists y)Rxy \vdash (\exists x)(\forall y)(\forall z)(Fx \rightarrow \sim Ryz)$
  5.  $(\forall x)(\forall y)(\forall z)(Rxy \rightarrow \sim Ryz) \vdash (\exists y)(\forall x)\sim Rxy$

## L2PL Natural Deduction Problem #1

Problem is :  $(\exists y)(\forall x)Rxy \vdash (\forall x)(\exists y)Rxy$

1	(1) $(\exists y)(\forall x)Rxy$	Premise
2	(2) $(\forall x)Rxb$	Ass ( $\exists E$ )
2	(3) $Rab$	2 $\forall E$
2	(4) $(\exists y)Ray$	3 $\exists I$
1	(5) $(\exists y)Ray$	1,2,4 $\exists E$
1	(6) $(\forall x)(\exists y)Rxy$	5 $\forall I$

## L2PL Natural Deduction Problem #2

Problem is :  $(\forall x)(\forall y)(\forall z)((Rxz \& Ryz) \rightarrow Rxy)$ ,  $(\forall x)Rxx \vdash (\forall x)(\forall y)(Rxy \rightarrow Ryx)$

1	(1) $(\forall x)(\forall y)(\forall z)((Rxz \& Ryz) \rightarrow Rxy)$	Premise
2	(2) $(\forall x)Rxx$	Premise
3	(3) $Rab$	Ass ( $\rightarrow$ I)
1	(4) $(\forall y)(\forall z)((Rbz \& Ryz) \rightarrow Rby)$	1 $\forall$ E
1	(5) $(\forall z)((Rbz \& Raz) \rightarrow Rba)$	4 $\forall$ E
1	(6) $(Rbb \& Rab) \rightarrow Rba$	5 $\forall$ E
2	(7) $Rbb$	2 $\forall$ E
2,3	(8) $Rbb \& Rab$	7,3 $\&$ I
1,2,3	(9) $Rba$	6,8 $\rightarrow$ E
1,2	(10) $Rab \rightarrow Rba$	3,9 $\rightarrow$ I
1,2	(11) $(\forall y)(Ray \rightarrow Rya)$	10 $\forall$ I
1,2	(12) $(\forall x)(\forall y)(Rxy \rightarrow Ryx)$	11 $\forall$ I

## L2PL Natural Deduction Problem #3

Problem is :  $(\forall x)(\forall y)(\forall z)((Rxz \& Ryz) \rightarrow Rxy)$ ,  $(\forall x)Rxx \vdash (\forall x)(\forall y)(\forall z)((Rxy \& Ryz) \rightarrow Rxz)$

1	(1) $(\forall x)(\forall y)(\forall z)((Rxz \& Ryz) \rightarrow Rxy)$	Premise
2	(2) $(\forall x)Rxx$	Premise
3	(3) $Rab \& Rbc$	Ass ( $\rightarrow$ I)
1	(4) $(\forall y)(\forall z)((Raz \& Ryz) \rightarrow Ray)$	1 $\forall$ E
1	(5) $(\forall z)((Raz \& Rcz) \rightarrow Rac)$	4 $\forall$ E
1	(6) $(Rab \& Rcb) \rightarrow Rac$	5 $\forall$ E
3	(7) $Rab$	3 $\&$ E
1,2	(8) $(\forall x)(\forall y)(Rxy \rightarrow Ryx)$	1,2 [Problem #2!]
1,2	(9) $(\forall y)(Rby \rightarrow Ryb)$	8 $\forall$ E
1,2	(10) $Rbc \rightarrow Rcb$	9 $\forall$ E
3	(11) $Rbc$	3 $\&$ E
1,2,3	(12) $Rcb$	10,11 $\rightarrow$ E
1,2,3	(13) $Rab \& Rcb$	7,12 $\&$ I
1,2,3	(14) $Rac$	6,13 $\rightarrow$ E
1,2	(15) $(Rab \& Rbc) \rightarrow Rac$	3,14 $\rightarrow$ I
1,2	(16) $(\forall z)((Rab \& Rbz) \rightarrow Raz)$	15 $\forall$ I
1,2	(17) $(\forall y)(\forall z)((Ray \& Ryz) \rightarrow Raz)$	16 $\forall$ I
1,2	(18) $(\forall x)(\forall y)(\forall z)((Rxy \& Ryz) \rightarrow Rxz)$	17 $\forall$ I

## L2PL Natural Deduction Problem #4

Problem is :  $(\forall x)Fx, \sim(\exists x)(\exists y)Rxy \vdash (\exists x)(\forall y)(\forall z)(Fx \rightarrow \sim Ryz)$

1	(1) $(\forall x)Fx$	Premise
2	(2) $\sim(\exists x)(\exists y)Rxy$	Premise
3	(3) $Fa$	Ass ( $\rightarrow$ I)
2	(4) $(\forall x)\sim(\exists y)Rxy$	2 SI (QS)
2	(5) $\sim(\exists y)Rby$	4 $\forall E$
2	(6) $(\forall y)\sim Rby$	5 SI (QS)
2	(7) $\sim Rbc$	6 $\forall E$
2	(8) $Fa \rightarrow \sim Rbc$	3,7 $\rightarrow$ I
2	(9) $(\forall z)(Fa \rightarrow \sim Rbz)$	8 $\forall$ I
2	(10) $(\forall y)(\forall z)(Fa \rightarrow \sim Ryz)$	9 $\forall$ I
2	(11) $(\exists x)(\forall y)(\forall z)(Fx \rightarrow \sim Ryz)$	10 $\exists$ I

## L2PL Natural Deduction Problem #5

Problem is :  $(\forall x)(\forall y)(\forall z)(Rxy \rightarrow \sim Ryz) \vdash (\exists y)(\forall x)\sim Rxy$

1	(1) $(\forall x)(\forall y)(\forall z)(Rxy \rightarrow \sim Ryz)$	Premise
2	(2) $\sim(\exists y)(\forall x)\sim Rxy$	Ass ( $\sim$ I)
2	(3) $(\forall y)\sim(\forall x)\sim Rxy$	2 SI (QS)
2	(4) $\sim(\forall x)\sim Rxa$	3 $\forall E$
2	(5) $(\exists x)\sim\sim Rxa$	4 SI (QS)
6	(6) $\sim\sim Rba$	Ass ( $\exists E$ )
2	(7) $\sim(\forall x)\sim Rxb$	3 $\forall E$
2	(8) $(\exists x)\sim\sim Rxb$	7 Taut.
9	(9) $\sim\sim Rcb$	Ass ( $\exists E$ )
1	(10) $(\forall y)(\forall z)(Rcy \rightarrow \sim Ryz)$	1 $\forall E$
1	(11) $(\forall z)(Rcb \rightarrow \sim Rbz)$	10 $\forall E$
1	(12) $Rcb \rightarrow \sim Rba$	11 $\forall E$
9	(13) $Rcb$	9 DN
1,9	(14) $\sim Rba$	12,13 $\rightarrow E$
6	(15) $Rba$	6 DN
1,6,9	(16) $\Delta$	14,15 $\sim E$
1,2,6	(17) $\Delta$	8,9,16 $\exists E$
1,2	(18) $\Delta$	5,6,17 $\exists E$
1	(19) $\sim\sim(\exists y)(\forall x)\sim Rxy$	2,18 $\sim I$
1	(20) $(\exists y)(\forall x)\sim Rxy$	19 DN

## Overview of the Course I

- Deductive Logic provides *formal theories of validity*. The deductive logician aims to *theoretically ground* our *informal* validity notion.
- In English, there are various argument forms or patterns that are intuitively or informally valid. We began with *sentential* forms like:  
Dr. Ruth is a man.  
(1) If Dr. Ruth is a man, then Dr. Ruth is 10 feet tall.  
∴ Dr. Ruth is 10 feet tall.
- Intuitively, (1)'s conclusion *follows-from* its premises. *If* the premises of (1) *were* all true, *then* (1)'s conclusion would also *have to be* true.
- Our first logical theory (LSL) correctly classifies this argument form (and many other valid English forms) as *valid*. A “success story” for LSL:

$p$ .  
(1<sub>LSL</sub>) If  $p$ , then  $q$ .  
∴  $q$ .

## Overview of the Course II

- However, there are many English arguments that are (intuitively, or “absolutely”) valid, but their LSL forms are *not* valid. For instance:

(2) Socrates is wise.  
 $\therefore$  Someone is wise.

- Intuitively, argument (2) is (“absolutely”) *valid*. But, if we try to translate this argument into LSL, we get the following *invalid* LSL form:

(2<sub>LSL</sub>)  $p.$   
 $\therefore q.$

- This is why we moved to the richer language LMPL, which subsumes LSL, and which adds additional structure that allows us to capture (2):

(2<sub>LMPL</sub>)  $Ws$   
 $\therefore (\exists x)Wx$

- Moreover, as we have seen, there are other valid English arguments that are beyond the reach of even the richer formal logical theory LMPL. *E.g.:*

(3) Brutus killed Caesar.  
 $\therefore$  Brutus killed someone and someone killed Caesar.

- If we were to symbolize this argument using LMPL, we would end-up with something like the following *invalid* LMPL argument form:

(3<sub>LMPL</sub>)  $Kb$   
 $\therefore (\exists x)Bx \ \& \ (\exists y)Ky$

Where  $Kx$ :  $x$  killed Caesar,  $Bx$ : Brutus killed  $x$ , and  $b$ : Brutus.

- The still richer language L2PL introduces 2-place predicates, such as  $Kxy$ :  $x$  killed  $y$ . With this relation in hand, we can capture (3) as:

(3<sub>L2PL</sub>)  $Kbc$   
 $\therefore (\exists x)Kbx \ \& \ (\exists y)Kyc$

- Of course, there are arguments beyond even L2PL's reach...

## Beyond L2PL I — Full First-Order Logic (LFOL)

- The full theory of first-order logic (LFOL) includes L2PL, plus  $n$ -place predicates, the identity relation  $=$ , and also function symbols.
- LFOL can capture even more valid arguments than L2PL. For instance, LFOL can capture arguments like the following mathematical one:

$$2 + 4 = 6$$

$$(4) \quad 3 \times 2 = 6$$

$$\therefore 2 + 4 = 3 \times 2$$

- Indeed, LFOL can capture just about any argument in just about any branch of modern mathematics. That's a lot of expressive power.
- In PHIL 140A, we study the full theory of first-order logic (LFOL). There, we give a semantics for LFOL, and we show that there is a sound and complete proof theory for LFOL (but, no decision procedure for  $\models$ !).
- Of course, even full first-order logic (LFOL) has its limitations...

## Beyond L2PL II — Second-Order Deductive Logic

- Some arguments involve quantification over not only objects but *properties*. These arguments are *second-order* and  $\therefore$  beyond LFOL.
- Leibniz (sometimes) talked as if the following argument were valid:  
(5)  $a$  and  $b$  have exactly the same (monadic) properties.  
 $\therefore a$  and  $b$  are identical.
- In second-order logic (SOL), (5) would be formalized as follows:  
(5<sub>SOL</sub>)  $(\forall P)(Pa \leftrightarrow Pb)$ .  
 $\therefore a = b$ .
- Note that the premise of (5) quantifies over (monadic) *predicates*.
- This is something that LFOL is not designed to do.
- We could also have an SOL which allows quantification over *relations*.
- Second-order logic is beyond 140A. It is touched upon (a little) in 140B.

## Beyond L2PL III — Non-Classical Deductive Logics

- All the logics I've mentioned are *classical* deductive logics. Not all logicians think classical logics capture our intuitive validity notions.
- Classical logics all share the following two properties:
  - (i) All arguments with contradictory premises (*e.g.*,  $p \ \& \ \sim p$ ) are valid.
  - (ii) All arguments with tautological conclusions (*e.g.*,  $p \ \vee \ \sim p$ ) are valid.
- Some logicians think (i) and/or (ii) are *counterexamples* to the classical theory of validity (as an explication of our informal “following-from”).
- Such logicians propose alternative formal theories of validity ( $\models^*$ ).
- Usually, non-classical logicians reject the classical (truth-functional) theory of the *conditional*. They adopt a non-classical conditional ( $\rightarrow^*$ ) which obeys constraints like the deduction theorem (relative to  $\models^*$ ).

$$p \models^* q \text{ if and only if } \models p \rightarrow^* q$$

- These and other fundamental philosophical questions about the nature of logic are addressed in our Philosophical Logic course (PHIL 142).

## Beyond L2PL IV — Inductive Logics

- Intuitively, not all “logically good” arguments are deductively valid. Some invalid arguments seem (intuitively) logically *better than* others:

(6)  $p$ . Someone is wise.  
 $\therefore q$ . Socrates is wise.

(7)  $r$ . Someone is either wise or unwise.  
 $\therefore q$ . Socrates is wise.

- *Inductive* logic should *theoretically ground* our intuition that (6) is a *logically stronger* argument than (7) is. Neither argument is *valid*.
- More ambitiously, an inductive logician might aim for a theory of “the *degree* to which the premises of an argument *confirm* its conclusion”.
- This ambitious project would aim to characterize a *function*  $c(\mathcal{C}, \mathcal{P})$ . And, an intuitive requirement would be that this function be such that:

$$c(q, p) > c(q, r)$$

- PHIL 148 is about probability and *inductive logic*. There, we explain how *probabilities* can be used to define various  $c$ -functions.

## Brief Review for Final

- The exam will contain 8 questions (same as the sample).
  - No L2PL interpretation *construction* problems on main exam.
  - There will also be an L2PL extra-credit problem. This problem will involve symbolization of an L2PL argument, and either interpretation construction or natural deduction.
  - Only try the extra-credit if you've got everything else nailed down.
- You will be given all natural deduction rules, but no truth-table definitions. The final rules handout is already posted online.
- You will be given 3 hours to complete the exam (you probably won't need it, but it's not a bad idea to go over things twice).
- Make sure to bring (several) blue books and a writing implement (preferably, a pencil). Turn in all materials at end.

# **Farewell**

Thanks for a great 6 weeks!

Good luck on your finals!

Have a great summer!