

---

# Examples from "A Decision Procedure for Probability Calculus with Applications"

Branden Fitelson  
Philosophy Department  
UC-Berkeley  
January 4, 2008

First, load the **PrSAT** package (freely downloadable from the **PrSAT** website: <http://fitelson.org/PrSAT/>).

```
<< PrSAT`
```

## ■ Example #1: A probability model in which three statements A, B, and C are *pariwise* independent but not *mutually* independent

Note: the additional equational side-constraints [ $\text{Pr}[A] = \frac{1}{10}$ ,  $\text{Pr}[B] = \frac{1}{10}$ ,  $\text{Pr}[C] = \frac{1}{10}$ ] are added here to reduce the number of variables of the problem, so that the model is found *much* more quickly. This is a very useful heuristic for speeding-up model searches. The option **Probabilities**→**Regular** guarantees that the model generated is *regular* (i.e., that it assigns nonzero probability to all state descriptions of the minimal sentential language required for the expression of the problem given). Finally, the option **BypassSearch**→**True** tells *Mathematica* to skip Blum's random search add-on, and send the problem straight to the decision procedure.

```
MODEL1 = PrSAT [
  {
    Pr[A ∧ B ∧ C] ≠ Pr[A] Pr[B] Pr[C],
    Pr[A ∧ B] == Pr[A] Pr[B],
    Pr[A ∧ C] == Pr[A] Pr[C],
    Pr[B ∧ C] == Pr[B] Pr[C],

    (* Heuristic -- add additional equational side-constraints *)
    Pr[A] ==  $\frac{1}{10}$ , Pr[B] ==  $\frac{1}{10}$ , Pr[C] ==  $\frac{1}{10}$ 
  },
  Probabilities → Regular,
  BypassSearch → True
]

{ {A → {a2, a5, a6, a8}, B → {a3, a5, a7, a8}, C → {a4, a6, a7, a8},
  Ω → {a1, a2, a3, a4, a5, a6, a7, a8}}, {a1 →  $\frac{1459}{2000}$ , a2 →  $\frac{161}{2000}$ ,
  a3 →  $\frac{161}{2000}$ , a4 →  $\frac{161}{2000}$ , a5 →  $\frac{19}{2000}$ , a6 →  $\frac{19}{2000}$ , a7 →  $\frac{19}{2000}$ , a8 →  $\frac{1}{2000}$ }}
```

**PrSAT** also includes a **TruthTable** function, which allows for the visualization of a model, as a stochastic truth-table:

**TruthTable[MODEL1]**

A	B	C	var	Pr
T	T	T	a <sub>8</sub>	$\frac{1}{2000}$
T	T	F	a <sub>5</sub>	$\frac{19}{2000}$
T	F	T	a <sub>6</sub>	$\frac{19}{2000}$
T	F	F	a <sub>2</sub>	$\frac{161}{2000}$
F	T	T	a <sub>7</sub>	$\frac{19}{2000}$
F	T	F	a <sub>3</sub>	$\frac{161}{2000}$
F	F	T	a <sub>4</sub>	$\frac{161}{2000}$
F	F	F	a <sub>1</sub>	$\frac{1459}{2000}$

If we set **BypassSearch**→**False**, then Blum's random-search add-on is consulted first, and a model is (usually) found (relatively) quickly *even without any side-constraints!*

**MODEL11 = PrSAT [**

```
{
  Pr[A ∧ B ∧ C] ≠ Pr[A] Pr[B] Pr[C],
  Pr[A ∧ B] == Pr[A] Pr[B],
  Pr[A ∧ C] == Pr[A] Pr[C],
  Pr[B ∧ C] == Pr[B] Pr[C]
},
```

**Probabilities** → **Regular**,

**BypassSearch** → **False**

**]**

{A → {a<sub>2</sub>, a<sub>5</sub>, a<sub>6</sub>, a<sub>8</sub>}, B → {a<sub>3</sub>, a<sub>5</sub>, a<sub>7</sub>, a<sub>8</sub>}, C → {a<sub>4</sub>, a<sub>6</sub>, a<sub>7</sub>, a<sub>8</sub>},

Ω → {a<sub>1</sub>, a<sub>2</sub>, a<sub>3</sub>, a<sub>4</sub>, a<sub>5</sub>, a<sub>6</sub>, a<sub>7</sub>, a<sub>8</sub>}}, {a<sub>1</sub> →  $\frac{84\,418 - 39\sqrt{4\,676\,097}}{56\,277}$ ,

a<sub>2</sub> →  $\frac{-42\,296 + 39\sqrt{4\,676\,097}}{168\,831}$ , a<sub>3</sub> →  $\frac{-42\,296 + 39\sqrt{4\,676\,097}}{168\,831}$ ,

a<sub>4</sub> →  $\frac{-42\,296 + 39\sqrt{4\,676\,097}}{168\,831}$ , a<sub>5</sub> →  $\frac{1}{999}$ , a<sub>6</sub> →  $\frac{1}{999}$ , a<sub>7</sub> →  $\frac{1}{999}$ , a<sub>8</sub> →  $\frac{42}{169}$ }}

TruthTable[MODEL11]

A	B	C	var	Pr
T	T	T	a <sub>8</sub>	$\frac{42}{169}$
T	T	F	a <sub>5</sub>	$\frac{1}{999}$
T	F	T	a <sub>6</sub>	$\frac{1}{999}$
T	F	F	a <sub>2</sub>	$\frac{-42\,296+39\sqrt{4\,676\,097}}{168\,831}$
F	T	T	a <sub>7</sub>	$\frac{1}{999}$
F	T	F	a <sub>3</sub>	$\frac{-42\,296+39\sqrt{4\,676\,097}}{168\,831}$
F	F	T	a <sub>4</sub>	$\frac{-42\,296+39\sqrt{4\,676\,097}}{168\,831}$
F	F	F	a <sub>1</sub>	$\frac{84\,418-39\sqrt{4\,676\,097}}{56\,277}$

We can also Evaluate probabilities on given models, as follows:

EvaluateProbability[{Pr[A | B], Pr[B | A ∧ ¬ C]}, MODEL11]

$$\left\{ \frac{42\,127}{168\,831 \left( \frac{42\,296}{168\,831} + \frac{-42\,296+39\sqrt{4\,676\,097}}{168\,831} \right)}, \frac{1}{999 \left( \frac{1}{999} + \frac{-42\,296+39\sqrt{4\,676\,097}}{168\,831} \right)} \right\}$$

% // N

{0.499521, 0.00400401}

### ■ A Simultaneous Countermodel to the **S**-instances of both (\*) and (†)

Again, the additional equational side-constraints [ $\Pr[H] = \frac{1}{2}$ ,  $\Pr[E1] = \frac{1}{4}$ ,  $\Pr[E2] = \frac{3}{4}$ ] are added to speed the model search, and **Probabilities→Regular** indicates that we are asking PrSAT to find a *regular* probability model.

```

MODEL2 = PrSAT [
  {
    Pr[H | E1] > Pr[H],
    Pr[H | E2] > Pr[H],
    Pr[H | E1] > Pr[H | E2],
    Pr[H | E1] - Pr[H | ¬E1] < Pr[H | E2] - Pr[H | ¬E2],
    Pr[H | E1 ∧ E2] - Pr[H | ¬E1 ∧ E2] == Pr[H | E1] - Pr[H | ¬E1],
    Pr[H | E2 ∧ E1] - Pr[H | ¬E2 ∧ E1] == Pr[H | E2] - Pr[H | ¬E2],
    Pr[H | E1 ∧ E2] - Pr[H | ¬(E1 ∧ E2)] < Pr[H | E2] - Pr[H | ¬E2],

    (* Heuristic -- add additional equational side-constraints *)
    Pr[H] ==  $\frac{1}{2}$ , Pr[E1] ==  $\frac{1}{4}$ , Pr[E2] ==  $\frac{3}{4}$ 
  },
  Probabilities → Regular,
  BypassSearch → True
]

```

```

{ {E1 → {a2, a5, a6, a8}, E2 → {a3, a5, a7, a8}, H → {a4, a6, a7, a8},
  Ω → {a1, a2, a3, a4, a5, a6, a7, a8}}, {a1 →  $\frac{147}{1024}$ , a2 →  $\frac{193}{5120}$ ,
  a3 →  $\frac{1341}{5120}$ , a4 →  $\frac{45}{1024}$ , a5 →  $\frac{291}{5120}$ , a6 →  $\frac{127}{5120}$ , a7 →  $\frac{1539}{5120}$ , a8 →  $\frac{669}{5120}$ }}

```

TruthTable[MODEL2]

E1	E2	H	var	Pr
T	T	T	a <sub>8</sub>	$\frac{669}{5120}$
T	T	F	a <sub>5</sub>	$\frac{291}{5120}$
T	F	T	a <sub>6</sub>	$\frac{127}{5120}$
T	F	F	a <sub>2</sub>	$\frac{193}{5120}$
F	T	T	a <sub>7</sub>	$\frac{1539}{5120}$
F	T	F	a <sub>3</sub>	$\frac{1341}{5120}$
F	F	T	a <sub>4</sub>	$\frac{45}{1024}$
F	F	F	a <sub>1</sub>	$\frac{147}{1024}$

## ■ A Simultaneous Countermodel to two claims concerning Hawthorne & Fitelson's new Bayesian approach to the raven paradox

The following single model shows that neither of the following two claims:

- (6)  $\Pr(H \mid \sim R \ \& \ \sim B) > \Pr(H)$   
 (7)  $\Pr(H \mid \sim R \ \& \ B) < \Pr(H)$

follows from the following three claims:

- (1)  $\Pr(R \mid H \ \& \ B) = 1$   
 (2)  $\Pr(\sim B) > \Pr(R)$   
 (C)  $\Pr(H \mid R) = \Pr(H \mid \sim B)$

Here, a regular model is impossible (since one of the constraints requires a zero probability for one of the state descriptions). But, by adding the constraint  $\Pr[(\sim H) \wedge B \wedge (\sim R)] > 0$ , we can ensure that this is the only zero in the model. And, as usual, we add equational side-constraints  $[\Pr[H] = \frac{60}{100}, \Pr[R] = \frac{20}{100}, \Pr[B] = \frac{10}{100}]$  to speed the model-finding process by reducing the number of free variables in the problem.

```

MODEL3 = PrSAT [
  {
    Pr[H ∧ R ∧ (¬ B)] == 0,
    Pr[(¬ H) ∧ B ∧ (¬ R)] > 0,
    Pr[¬ B] > Pr[R],
    Pr[H | R] == Pr[H | ¬ B],
    Pr[H | (¬ R) ∧ (¬ B)] < Pr[H],
    Pr[H | B ∧ (¬ R)] > Pr[H]
  },
  Constraints → {Pr[B] == 1/10, Pr[H] == 1/2, Pr[R] == 1/10},
  BypassSearch → True
]

{B → {a2, a5, a6, a8}, H → {a3, a5, a7, a8},
 R → {a4, a6, a7, a8}, Ω → {a1, a2, a3, a4, a5, a6, a7, a8}},
 {a1 → 141/320, a2 → 21/2560, a3 → 225/512, a4 → 51/2560, a5 → 3/256, a6 → 1/32, a7 → 0, a8 → 25/512}

```

TruthTable[MODEL3]

B	H	R	var	Pr
T	T	T	a <sub>8</sub>	$\frac{25}{512}$
T	T	F	a <sub>5</sub>	$\frac{3}{256}$
T	F	T	a <sub>6</sub>	$\frac{1}{32}$
T	F	F	a <sub>2</sub>	$\frac{21}{2560}$
F	T	T	a <sub>7</sub>	0
F	T	F	a <sub>3</sub>	$\frac{225}{512}$
F	F	T	a <sub>4</sub>	$\frac{51}{2560}$
F	F	F	a <sub>1</sub>	$\frac{141}{320}$

We can also solve this one (quickly) with Blum's random search add-on, and with no side-constraints:

```
PrSAT [
  {
    Pr[H ∧ R ∧ (¬ B)] == 0,
    Pr[(¬ H) ∧ B ∧ (¬ R)] > 0,
    Pr[¬ B] > Pr[R],
    Pr[H | R] == Pr[H | ¬ B],
    Pr[H | (¬ R) ∧ (¬ B)] < Pr[H],
    Pr[H | B ∧ (¬ R)] > Pr[H]
  },
  BypassSearch → False
]

{ {B → {a2, a5, a6, a8}, H → {a3, a5, a7, a8},
  R → {a4, a6, a7, a8}, Ω → {a1, a2, a3, a4, a5, a6, a7, a8}},
  {a1 →  $\frac{36\,693}{94\,240}$ , a2 →  $\frac{2}{15}$ , a3 → 0, a4 →  $\frac{1}{93}$ , a5 →  $\frac{11}{32}$ , a6 →  $\frac{7}{57}$ , a7 → 0, a8 → 0}}
```

## ■ Theorems and Countermodels from Sobel's "Lotteries and Miracles"

Since Sobel's problems only involve two atomic sentences, no heuristics are needed to yield a fast solution by the decision procedure (and Blum's random search add-on is also not necessary, since the decision procedure is quite fast in such cases).

### ■ A PrSAT model showing that Sobel's (1)–(3) do *not* entail (4)

```
MODEL4 = PrSAT [
  {
    Pr[T] <  $\frac{1}{2}$ ,
    Pr[T | W] >  $\frac{1}{2}$ ,
    Pr[W | T] >  $\frac{1}{2}$ ,
    Pr[T | ¬ W] > Pr[W]
  },
  Probabilities → Regular,
  BypassSearch → True
]

{ {T → {a2, a4}, W → {a3, a4}, Ω → {a1, a2, a3, a4}},
  {a1 →  $\frac{6033}{8192}$ , a2 →  $\frac{1007}{8192}$ , a3 →  $\frac{1}{64}$ , a4 →  $\frac{1}{8}$ }}
```

TruthTable[MODEL4]

T	W	var	Pr
T	T	a <sub>4</sub>	$\frac{1}{8}$
T	F	a <sub>2</sub>	$\frac{1007}{8192}$
F	T	a <sub>3</sub>	$\frac{1}{64}$
F	F	a <sub>1</sub>	$\frac{6033}{8192}$

A PrSAT model showing that Sobel's (1)–(3) do *not* entail (5)

```

MODEL5 = PrSAT [
  {Pr [T] < 1/2,
   Pr [T | W] > 1/2,
   Pr [W | T] > 1/2,
   Pr [T | W] - Pr [T | ¬ W] < Pr [¬ W] - Pr [W]
  },
  Probabilities → Regular,
  BypassSearch → True
]

{ {T → {a2, a4}, W → {a3, a4}, Ω → {a1, a2, a3, a4}},
  {a1 → 51/64, a2 → 1/16, a3 → 1/16, a4 → 5/64} }

```

TruthTable[MODEL5]

T	W	var	Pr
T	T	a <sub>4</sub>	5/64
T	F	a <sub>2</sub>	1/16
F	T	a <sub>3</sub>	1/16
F	F	a <sub>1</sub>	51/64

■ A PrSAT "Proof" that Sobel's (1)–(3) do entail the *disjunction* (4) ∨ (5)

```

PrSAT [
  {Pr [T] < 1/2,
   Pr [T | W] > 1/2,
   Pr [W | T] > 1/2,
   Not [Or [Pr [T | ¬ W] < Pr [W], Pr [T | W] - Pr [T | ¬ W] > Pr [¬ W] - Pr [W]]]
  },
  BypassSearch → True
]

{}

```