

POLLOCK ON PROBABILITY IN EPISTEMOLOGY

BRANDEN FITELSON

ABSTRACT. John Pollock has done a lot of interesting and important work on the metaphysics and epistemology of probability over several decades. In *Thinking About Acting* [14], we find many fascinating and thought provoking ideas and arguments (both old and new) about probability. Owing to limitations of space, I will be confining my remarks to a handful of issues addressed in [14] pertaining to probability, logic, and epistemology. First, I will discuss some of Pollock’s arguments against Bayesian Epistemology (BE). Here, I’ll try to defend (BE) from what I take to be less than decisive objections. Then, I will make some critical remarks concerning Pollock’s alternative approach to “probabilistic epistemology”, which is based on his (non-Bayesian) theory of “nomic probability” [13].¹

1. SOME REMARKS ON POLLOCK’S CRITIQUE OF BAYESIAN EPISTEMOLOGY

In chapter 6, Pollock offers various criticisms of (BE). Before delving into some of Pollock’s criticisms, I would like to begin by pointing out some puzzling things Pollock says about the probability calculus. This will lead, naturally, into a more substantive discussion of Pollock’s critique of (BE). According to Pollock [14, p. 83], the probability calculus can be axiomatized as follows:

- (1) $\mathbf{PROB}(P \ \& \ \sim P) = 0$.
- (2) $\mathbf{PROB}(P \vee \sim P) = 1$.
- (3) $\mathbf{PROB}(P \vee Q) = \mathbf{PROB}(P) + \mathbf{PROB}(Q) - \mathbf{PROB}(P \ \& \ Q)$.
- (4) If P and Q are logically equivalent, then $\mathbf{PROB}(P) = \mathbf{PROB}(Q)$.

Pollock claims that the above axiomatization is equivalent to Kolmogorov’s *probability calculus* (PC) [10].¹ In fact, Pollock’s intended interpretation of his axiomatization — which I’ll call (PC’), for short — is *logically incomparable* to (PC). That is, (i) there are some theorems of (PC) that are not theorems of (PC’), and (ii) there are some theorems of (PC’) that are not theorems of (PC). Let’s take (ii) first. The probability calculus defines probability functions $\mathbf{PROB}(\cdot)$ over *sentential languages* \mathcal{L} . As such, if (PC’) is to be equivalent to (PC), then the (schematic) axioms (1)–(4) must be *relativized* to some such \mathcal{L} . That is, the metavariables “ P ” and “ Q ” in Pollock’s (1)–(4) must be understood as *ranging over sentences of some sentential language* \mathcal{L} . If we do not do this, then we may falsely interpret “logically equivalent” in (4) as something stronger than “*tautologically equivalent in* \mathcal{L} ,” which is all “logically equivalent” means in (PC). [As we’ll see shortly, Pollock’s critique of (BE) makes use of just such a stronger reading of the locution “logically equivalent” in (4).] As a result, (PC’) contains “theorems” that are *not* theorems of (PC). For instance, Pollock’s (PC’) will entail that $\mathbf{PROB}(P) = \mathbf{PROB}(Q)$, for many P and Q that are

Date: 07/02/09.

¹Strictly speaking, Kolmogorov gives a set-theoretic, and not a logical axiomatization of (PC). But, one can give an (extensionally) equivalent logical axiomatization. See [5, §1] for an axiomatization of (PC) that is along these lines.

not even *expressible* in any sentential language \mathcal{L} (e.g., first-order or higher-order equivalences). I’ll return to this, below, in my discussion of Pollock’s critique of (BE). But, first, let me illustrate (i). The following is an axiom of Kolmogorovian (PC) [5]:

- (5) For all $P \in \mathcal{L}$, $\mathbf{PROB}(P) \geq 0$.

Unfortunately, (5) is *not* a theorem of (PC’) — even when it is (properly) restricted to sentential languages \mathcal{L} . To see this, we can construct a simple counterexample to (5) in a properly \mathcal{L} -relativized version of (PC’). Let \mathcal{L} contain just one atomic sentence A . Hence, only four distinct propositions can be expressed in \mathcal{L} : A , $\sim A$, $A \ \& \ \sim A$, and $A \vee \sim A$. Now, let $\mathbf{PROB}(\cdot)$ be defined on \mathcal{L} , as follows:

- (6) $\mathbf{PROB}(A) = 2$
- (7) $\mathbf{PROB}(\sim A) = -1$
- (8) $\mathbf{PROB}(A \vee \sim A) = 1$
- (9) $\mathbf{PROB}(A \ \& \ \sim A) = 0$

This $\langle \mathcal{L}, \mathbf{PROB} \rangle$ pair satisfies all of Pollock’s (PC’) axioms (1)–(4), but it also violates Kolmogorov’s (5), since $\mathbf{PROB}(\sim A) = -1 < 0$.² Therefore, Pollock’s (PC’) is *both too strong* [(ii)] *and too weak* [(i)] to be a proper candidate for an equivalent formulation of (PC). Problem (i) is easily fixed, by adding (5) as an axiom to a properly \mathcal{L} -relativized rendition of (PC’). But, problem (ii) is deeper and more intertwined with Pollock’s thinking about Bayesianism. If we fix problem (ii) by limiting the axioms of (PC’) to sentential languages \mathcal{L} — and we bear this limitation in mind when we apply (PC’) to (BE) — then some of Pollock’s central criticisms of (BE) will be threatened. Allow me to explain.

One of Pollock’s main lines of criticism of (BE) is that it entails a kind of *logical omniscience*. As he rightly points out [14, p. 94], some necessary truths seem (intuitively) *unjustified* for some epistemically rational agents (S). But, Pollock claims, Bayesian epistemology cannot make sense of this, if it is to use $\mathbf{PROB}(p)$ as a way of gauging the *degree to which* p *is justified* (for S). He says:

If Q is a necessary truth, it is logically equivalent to $(P \vee \sim P)$, so it follows from axioms (2) and (4) that every necessary truth has a \mathbf{PROB} of 1.

I think this is highly uncharitable to the Bayesian epistemologist. First, this rests on a misunderstanding of (PC), which *only* entails that *tautologies of* \mathcal{L} must be assigned a \mathbf{PROB} of 1. Second, it rests on an implausible assumption about “necessary truths” — that they are all logically equivalent to the simple tautology $P \vee \sim P$. I’m not sure what Pollock has in mind here, but I don’t see why a Bayesian (or anyone else) should be saddled with such a strong commitment. As a result, it’s unclear what reason Bayesians could have for insisting that all necessary truths be assigned the same probability as a tautology. It seems to me that there are better ways to think about (PC) and (BE).

Garber [6] explains how (PC) — when properly construed and applied — can be used by Bayesian epistemologists to model logically non-omniscient agents. And, he uses his approach to give a novel and compelling resolution of the so-called “old

²Pollock is in good company here. Skyrms’s [17, ch. 6] axiomatization has exactly the same deficiency. I owe this counterexample to Skyrms’s (and Pollock’s) theory to Mike Titelbaum. As Carnap [3, p. 341] notes, it is surprisingly easy to give equivalent-*looking* axioms for (PC), which are *non-equivalent*. This happens a lot in the literature on (PC).

evidence problem” of Bayesian confirmation theory (a branch of Bayesian epistemology). I won’t get into the details of Garber’s approach here. But, I’ll give the basic idea behind it. Pollock’s criticism presupposes a *very rich* notion of “logical equivalence” in his interpretation of (PC). As I have explained, however, (PC) has an *impoverished* notion of “logical equivalence” — *tautological equivalence in some sentential \mathcal{L}* [5]. While this impoverishment may seem like a shortcoming³ — it can be a virtue. It allows Bayesian epistemologists to model agents who may *only* be omniscient about the tautologies of some sentential language \mathcal{L} . By exploiting the fact that atomic sentences are not tautologically related to each other, we can then use this “impoverishment” to model *ignorance* of “higher” logical truths, which are not expressible in \mathcal{L} . Following Garber, we can do so by *extra-systematically interpreting* the atomic sentences of \mathcal{L} . For instance, we could have a language with three atomic sentences A , B , and C , where “ C ” gets extra-systematically interpreted as “ A entails B ”, and where this “entailment” is (say) first-order (but not sentential). Then, we could add extra-systematic probabilistic constraints to our probability model, which would *selectively capture* such “higher” logical knowledge on the part of the agent being modeled. For example, by adding the following extra-systematic constraint, we can model an agent who knows that “*modus ponens*” for “entails” is *extra-systematically valid* (in this instance):

$$(10) \text{PROB}(B | A \ \& \ C) = 1.$$

We could also allow this conditional **PROB** to be less than 1, in which case we’d be modeling an agent who is *ignorant* of this “extra-systematic *modus ponens*”. In this way, we can model agents who are justified in believing some extra-systematic (logical, conceptual, or other) necessary truths, but not others. And, that gives a sophisticated Bayesian epistemologist the wherewithal to overcome this criticism of Pollock. Of course, Garber’s framework still presupposes *some* logical omniscience, and this leaves the Bayesian vulnerable to some objections. Indeed, Pollock [14, p. 94] rightly points out that sometimes people aren’t even justified in believing some tautologies in simple languages \mathcal{L} . And, that problem will still plague even a Garberian approach to Bayesian epistemology.⁴ However, as Pollock himself notes [14, p. 95], tautologies are always *warranted*. So, presumably, Pollock’s logical omniscience objection would not undermine a Garberian application of **PROB** to the modeling of degrees of *warrant*. In any case, Pollock has another objection to this sort of Bayesian epistemology.

According to Pollock, Bayesian epistemologists who claim that **PROB** is a measure of degree of warrant (or justification) are unable to explain the role of *reasoning* in epistemology. Pollock seems to think that the following is a *desideratum* for any adequate (formal) epistemology [14, p. 95]:

- (11) Any adequate (formal) epistemology must be able to explain why deductive inference from multiple uncertain premises can be expected to *preserve justification (and/or warrant)*.

Pollock argues that (BE) is unable to meet requirement (11). He says [14, p. 97]:

If degrees of warrant satisfy the probability calculus, then ... we can only be confident that a deductive argument takes us from warranted premises to a warranted conclusion if all the inferences are probabilistically valid.

Where, an inference of the form $P_1, \dots, P_n \therefore Q$ is *probabilistically valid* just in case its conclusion Q is at least as probable as its least probable premise — that is, iff for all i : **PROB**(Q) \geq **PROB**(P_i). As it turns out, no deductively valid form of inference with more than one premise is probabilistically valid *in this sense*. That explains why Pollock thinks Bayesian epistemology cannot satisfy (11). The reason Pollock thinks violating (11) is undesirable is that he thinks violating (11) prevents probabilism from being able to explain how we can reason “blindly” from multiple warranted (or justified) premises, using a deductively valid inference, and expect that the conclusion will also be warranted (or justified). Since “blind deductive reasoning” seems integral to epistemology, this would be a serious shortcoming of (BE) — or, more generally, of any *probabilistic* epistemology.

Strictly speaking, it is true that Bayesianism *so construed* can’t satisfy (11) *in this sense*. But, I wonder why one would want to *both* construe Bayesian epistemology in this way, *and* understand “probabilistic validity” in this way. It seems clear to me that many contemporary Bayesian epistemologists would *neither* want to equate **PROB** and *degree of warrant* (or *degree of justification*, for that matter) *nor* explicate *probabilistic validity* in the way Pollock proposes. Let’s take the second point first. There is quite a long tradition of what is known as *probability logic* (PL). In recent years, probability-logicians like Adams [1] and Hailperin [7] have done a great deal of work on various notions of “probabilistic validity”. Two important points about (PL) are in order here. First, the notion of “probabilistic validity” that is typically used in (PL) circles is not the one Pollock has in mind. Adams [1, p. 57] defines a different notion, which I will call **PROB-validity**. I won’t give his definition of **PROB-validity** here, but I will discuss one important consequence of the definition, just to give a sense of how it differs from Pollock’s “probabilistic validity”. Let $\mathbf{U}(p) = \underline{\underline{1}} - \text{PROB}(p)$ be the *uncertainty* of p . And, consider an inference of the form $P_1, \dots, P_n \therefore Q$. Such an inference will be **PROB-valid** in Adams’s sense *only if*⁵ the uncertainty of the conclusion is no greater than the sum of the uncertainties of the premises — that is, *only if* $\mathbf{U}(Q) \leq \sum_{i=1}^n \mathbf{U}(P_i)$. In other words, the uncertainty of the conclusion of a **PROB-valid** inference *will never exceed the sum-total of the uncertainties of its premises*. Moreover, it is a fundamental theorem of (PL) that *all deductively valid arguments are **PROB-valid***. So, in this sense, a Bayesian (probabilist) who adopts Adams’s notion of **PROB-validity**, can explain why (in one precise sense) conclusions of deductively valid inferences will never be *more unwarranted* (or *more unjustified*) than the premises already were. Of course, this presupposes a different epistemic *explanandum* than Pollock has in mind in (11). But, in the interest of giving (PC) and (BE) a fair hearing, it is worth noting that other notions of

³A bit later in the text, Pollock discusses a related logical impoverishment of (PC), and he complains that it is a shortcoming. On page 108, Pollock rightly points out that (PC) does not say anything (systematically) about probabilities over *open* first-order sentences. This is true, of course. But, something much stronger is true — namely that (PC) doesn’t say anything (systematically) about **PROB**abilities over *anything other than sentential* languages \mathcal{L} .

⁴Having conceded this point, it is worth mentioning that this problem is far less pressing than the problem Pollock has in mind — which would saddle proponents of (BE) with the commitment to assign probability 1 to *all necessary truths*. The main point I want to get across here is that proponents of (BE) have the theoretical tools to distinguish various “levels” of ideal epistemic rationality. As such, their framework is not as hopeless as Pollock makes it sound.

⁵This is only a necessary condition for **PROB-validity**, which is why it is not suitable as a definition [1, p. 57].

“probabilistic validity” have been investigated by people who are interested in just the sort of deductive inferences from multiple uncertain premises that Pollock is talking about. Putting these alternative (PL)-investigations of “uncertain deductive inference” to one side, I want to make a second point about (PL) — that it can be illuminating, even with respect to Pollock’s *explanandum* [(11)].

Consider *modus ponens* for material implication “ \supset ” (“ \supset -MP”, for short). This is a very common and important multi-premise deductive inference that (I take it) is used often in the sort of “blind” deductive reasoning Pollock has in mind. Given Pollock’s definition, \supset -MP is not “probabilistically valid”. But, (PL) allows us to be more precise in our “diagnosis”. Here is a (PL)-fact about \supset -MP:

(12) If $\mathbf{PROB}(P) > 1 - \epsilon$ and $\mathbf{PROB}(P \supset Q) > 1 - \epsilon$, then $\mathbf{PROB}(Q) > 1 - 2\epsilon$.⁶

This is a classical theorem of (PL) [7, p. 205]. While (12) entails that *some* degree of \mathbf{PROB} can be “lost” in (material) *modus ponens* inferences, it also tells us that, when the premises are *highly* probable, the *amount* of \mathbf{PROB} that can be lost in \supset -MP is rather small. Now, imagine a Bayesian epistemologist who wants to defend the claim that degree of justification \mathbf{DJ} (or degree of warrant \mathbf{DW}) is a \mathbf{PROB} . I don’t think someone like this is at a *complete* loss to explain (even on Pollock’s terms) how \supset -MP can *often* be “blindly” applied, *while preserving justified-ness* (or warranted-ness). Let us (naïvely) assume the following \mathbf{PROB} -reduction of justified-ness (or warranted-ness):

(13) S is justified (warranted) in believing p iff $\mathbf{PROB}(p) > 1 - 2\epsilon$, for some suitably “small” ϵ ; and, S is *highly* justified (warranted) in believing p iff $\mathbf{PROB}(p) > 1 - \epsilon$.⁷

In light of (12), a Bayesian who endorses (13) *can* explain how we may “blindly” do \supset -MP — in cases where the premises are all *highly* justified, since (12) entails that if the premises of a \supset -MP-inference are all highly justified, then the conclusion must be justified. Granted, this isn’t *as general* an explanation of “blind \supset -MP” as a Bayesian would have if \supset -MP were “probabilistically valid” in Pollock’s sense. But, I don’t see why this isn’t explanatory *at all* — even with respect to Pollock’s *explanandum* (or an *explanandum* that is very similar to Pollock’s). A similar strategy can be employed for \supset -transitivity, in light of the following classical theorem of (PL) [7, p. 205]:

(14) If $\mathbf{PROB}(P \supset Q) > 1 - \epsilon$ and $\mathbf{PROB}(Q \supset R) > 1 - \epsilon$, then $\mathbf{PROB}(P \supset R) > 1 - 2\epsilon$.

Furthermore, if we talk about *indicative modus ponens* (\rightarrow -MP), rather than *material modus ponens* (\supset -MP), then things get even more interesting.⁸ Many people [1, 2, 4]

⁶I haven’t said anything yet about the *interpretation* of \mathbf{PROB} . This is intentional. It seems to me that Pollock’s objections are not restricted to (say) *subjective* (BE). Rather, he’s taking on just about *any* kind of \mathbf{PROB} abilistic reduction of \mathbf{DJ} or \mathbf{DW} . I presume this would include *non*-subjective probabilists about evidential support, such as Carnap [3], Williamson [19], and Keynes [9], as well as *subjective* (BE)-ers, such as Skyrms [17], Joyce [8], and others. I’ll return to this issue in section two, below. But, in the meantime, I will assume that \mathbf{PROB} is whatever probability function a particular advocate of (BE) has in mind. This will vary, but in a way that is orthogonal to this line of Pollock’s objections.

⁷Of course, I do *not* mean to endorse (13), *nor* do I mean to saddle the proponent of (BE) with it. I am only introducing it here for dialectical purposes — to bring out what I think is an exaggeration in Pollock’s objection to (BE).

⁸Various commentators have recently come to the view that \rightarrow -MP isn’t even deductively valid [12, 11]. I will put that controversy to one side here, and I will suppose that *modus ponens* is deductively

think that the probability of the indicative conditional $P \rightarrow Q$ goes according to the conditional probability $\mathbf{PROB}(Q | P)$. If that’s right, then we get an even better result of (PL) for the dialectical purposes at hand, namely:

(15) If $\mathbf{PROB}(P) > 1 - \epsilon$ and $\mathbf{PROB}(P \rightarrow Q) = \mathbf{PROB}(Q | P) > 1 - \epsilon$, then $\mathbf{PROB}(Q) > (1 - \epsilon)^2$.

That is, supposing that \rightarrow -MP is valid and that the equation $\mathbf{PROB}(P \rightarrow Q) = \mathbf{PROB}(Q | P)$ is correct, this means that even less \mathbf{PROB} can be “lost” in \rightarrow -MP inferences than in \supset -MP inferences. And, so, the analogous Bayesian strategy is even more explanatory (in Pollock’s sense) in that case. There are limits to this strategy, since some multi-premise deductive arguments won’t even be guaranteed to preserve warrant/justification in cases where all the premises are *highly* warranted/justified. But, there is a fully general theory of “probability logic”, which furnishes such results for many classically deductively valid argument forms [7]. To my mind, this (to some extent) softens the impact of Pollock’s objection to thinking of degree of justification or degree of warrant as a \mathbf{PROB} .

Finally, I want to return to the question of whether Bayesians *should* (or *do*) think of degree of justification (or warrant) as a \mathbf{PROB} -function. I think many contemporary Bayesians would *not* want to do this, but for reasons that are independent of the considerations we just discussed in connection with Pollock’s objection. Bayesian epistemologists typically distinguish two types of “evidential support” or “confirmation” — *firmness* and *increase in firmness* [3, new preface]:

- **Confirmation as firmness.** E confirms _{f} H , relative to background evidence K if and only if $\mathbf{PROB}(H | E \ \& \ K) > t$, for some threshold value t (typically, $t > 1/2$).
- **Confirmation as increase in firmness.** E confirms _{i} H , relative to background evidence K if and only if $\mathbf{PROB}(H | E \ \& \ K) > \mathbf{PROB}(H | K)$.

The firmness concept is sometimes called “absolute” confirmation, and the increase in firmness concept is sometimes called “incremental” confirmation. The distinction between them is just the distinction between *high conditional probability* vs *probabilistic relevance*. While these two concepts are closely related to each other, they can come apart in some rather important ways. Here is an example (to which I’ll return in section 2, below) that illustrates the confirms _{f} /confirms _{i} distinction.

The example involves a 35-year old American man named Jim, who has received a positive test result for a rare disease X . Only 1 in 10,000 35-year-old males in the U.S. has disease X . But, the test for X is very highly reliable — it has very low false-positive and false-negative rates (each of these error rates is 1/1000). That is, if you have disease X , then there is only a 1/1000 chance of a false negative from an X -test, and if you don’t have X , then there is only a 1/1000 chance of a false positive from an X -test. Let a denote Jim, let Nx assert that x does *not* have disease X , and let Px assert that x has received a (single) positive test result for disease X . In this case, we (intuitively) have the following probabilistic facts, where

valid for the indicative conditional. But, it is worth noting that, if these commentators are right, then “blind deductive \rightarrow -MP reasoning” would not be kosher. I think that would undermine Pollock’s dialectical position vis-a-vis (BE). But, I can’t go into that here.

K is the background evidence contained in the above story about Jim, the disease, and the test (and **PROB** may be interpreted in various ways⁹):

- **PROB**($Na \mid Pa \ \& \ K$) is *high* (specifically, it's approximately 9/10).
- **PROB**($Na \mid Pa \ \& \ K$) is *significantly less than* **PROB**($Na \mid K$).

In other words, Pa confirms _{f} Na , relative to K ; but Pa disconfirms _{i} Na , relative to K . Question: are we justified/warranted in *believing* Na on the basis of Pa (given background knowledge K)? On the one hand (the firmness hand), Na is *highly probable*, given Pa (and K). On the other hand (the increase in firmness hand), Pa is *strongly negatively relevant* to the probability of Na (given K). This conflict between confirms _{f} and disconfirms _{i} seems to pull intuitions about *believing* Na on the basis of Pa in opposite directions. Many advocates of (BE) seem to endorse the following.¹⁰

- (16) A *necessary condition* for being justified/warranted in believing H on the basis of E (given background evidence/knowledge K) is that E *does not disconfirm* _{i} H , relative to K .

If (16) is correct, then we would *not* be warranted/justified in believing Na on the basis of Pa (given background knowledge K) — *despite* the fact that **PROB**($Na \mid Pa \ \& \ K$) is high.¹¹ This sort of consideration is a reason for many advocates of (BE) to reject the idea that the degree to which E justifies/warrants H (relative to background knowledge K) is **PROB**($H \mid E \ \& \ K$). And, this consideration is orthogonal to the considerations raised by Pollock's objections concerning "blind (uncertain) deductive reasoning". This example also provides a nice *segué* into section two, where I will appeal to similar considerations to pose a challenge to Pollock's alternative "probabilistic epistemology".

2. SOME WORRIES ABOUT POLLOCK'S ALTERNATIVE "PROBABILISTIC EPISTEMOLOGY"

Pollock rejects (BE), but he still thinks that probabilities (of some kind) are important in epistemology. Pollock's alternative is what I will call a theory of *defeasible probabilistic reasoning* (DPR). Pollock's (DPR) has three main components, each of which differs in important ways from (BE).

The first component of Pollock's (DPR) involves *indefinite* probabilities. The probability calculus (and the example we discussed above) involves only *definite* probabilities — probabilities over *closed* sentences (*i.e.*, *propositions*). Pollock's (DPR) theory involves *nominal probability* [13] functions **prob**, which (formally) take *open* sentences as arguments. For instance, **prob**($Nx \mid Px$) is meaningful in Pollock's theory, and it denotes "the proportion of physically possible P 's that would

⁹As I explained in footnote 6, I am remaining as neutral as possible on the *interpretation* of **PROB** here. I will return to this issue in section two, below. In this example, I think the probabilistic "facts" I cite are robust across various interpretations of **PROB**. And, I think I'm not doing any harm here to Pollock's usage of **PROB** for *definite* probabilities.

¹⁰Roger White [18, §5] seems to assume something like (16) in his Bayesian criticism of epistemic dogmatism. Tim Williamson [19, chs. 9 & 10] seems to require some *probabilistic relevance* in his account of "justification". And, Tomoji Shogenji [16] defends a precise, probabilistic theory of **DJ**, according to which **DJ** is *not* a confirms _{f} -function (*i.e.*, not a conditional **PROB** function), but rather a confirms _{i} -function. I'm inclined to think that that a proper Bayesian theory of **DJ** (if there be such) will have to be sensitive to *both* firmness *and* increase in firmness considerations.

¹¹Note that we can make **PROB**($Na \mid Pa \ \& \ K$) as high as we like, just by fiddling with the numbers given in K .

be N 's". So, Pollock is talking about a kind of *objective, physical* probability, which is *indefinite*. This differs from the **PROB**'s of (BE) in several respects. First, the **PROB**'s of (BE) are (in some sense) *epistemic* probabilities. And, while there is disagreement among advocates of (BE) as to whether epistemic probabilities are subjective or objective (see footnote 6), it is clear that **PROB**'s are *not physical* probabilities. Second, Pollock's **prob**'s are indefinite, while (BE)'s **PROB**'s are definite. This is also important, since both Pollock and the advocates of (BE) want to make inferences *about particulars*. Pollock will do this *via defeasible reasoning* from his indefinite, nomic **prob**'s (plus definite statements about particulars) to (other) definite statements about particulars. Bayesians will do this *via* direct appeals to definite **PROB**abalistic "facts". Finally, Pollock's indefinite **probabilities** *formally* differ from (PC)'s **PROB**'s in various ways. Pollock has developed a sophisticated formal theory of **prob**, as well as some ingenious computer programs for calculating and proving general claims about **prob**'s. Unfortunately, I don't have the space to discuss any of that formal work here.¹² Next, I will illustrate how Pollock's (DPR) approach differs from (BE) on our example above. But, first, I need to mention the other two components of Pollock's theory of defeasible probabilistic reasoning.

The second component of Pollock's (DPR) will require some account of *how we can come to know* the (true) values of (or, at least, ranges of values of or inequalities involving) salient nomic probabilities. Among other things, this will have to give us some grip on how we might come to know something about the "true proportionality function ρ over nomologically possible worlds". I put this locution in quotation marks, because I am rather skeptical that there *are* such proportionality functions, and/or that we can come to *know* what they are. But, because my space is limited here, I won't be able to get into the (rather extensive) metaphysical and epistemological worries I have about "proportions of nomologically possible worlds"-talk. Pollock does have a lot to say about this second component. And, I refer the interested reader to his 1990 book on nomic probability [13].

The third component (and the most salient one for my purposes here) of Pollock's (DPR) involves *principles of defeasible reasoning* from facts about particulars + facts about nomic probabilities to further facts about particulars. Pollock discusses various principles of this kind throughout his work on nomic probability. I will focus on just one of these, which is central to his approach — the so-called *statistical syllogism* (SS). Pollock gives various formulations of (SS) in his work. I will use the following formulation from the book [14, p. 235], which is most convenient for my purposes:

- (SS) If F is projectible with respect to G and $r > 0.5$, then " $Gc \ \& \ \mathbf{prob}(Fx \mid Gx) \geq r$ " is a defeasible reason for believing " Fc ", the strength of the reason depending upon the value of r .

Let's apply (SS) to our example above, where we will just take the description of Jim, the disease, and the test as part of our background knowledge. Then, I presume, we have a case in which we ought to be able to apply (SS). First, I presume that our background knowledge will (somehow) allow us to know the salient nomic

¹²Pollock has made a lot of progress on the formal/computational side of his theory since the book was written. I have had the pleasure of reading a more recent manuscript [15], which develops the formal side in much more detail and generality. I have also benefited from a very edifying email correspondence with John about his quite extensive and impressive computational work on **prob**, and its relation to my recent computational work on **PROB** [5].

probability $\mathbf{prob}(Nx | Px)$. Specifically, I presume (and hope) it turns out that we can know that $\mathbf{prob}(Nx | Px) > 0.5$. Indeed, I presume (and hope) it turns out that we can know this nomic probability is around 0.9 in the example at hand. Then, it seems to me that (SS) should imply the following, in our example (since I take it we have projectibility here as well):

- (17) Pa is a defeasible reason to believe Na (given what we know about the example in question). Moreover, Pa is a *strong* (defeasible) reason to believe Na (and we can make it *as strong a reason as we like*, just by turning-up the numbers in our background story about the case).

On its face, (17) suggests that a positive test result from a highly reliable test provides an (arbitrarily) strong reason to believe that the disease is *absent!* I find that counter-intuitive. And, I think the story that advocates of (BE) tell about confirms_f vs confirms_i furnishes a pretty plausible explanation of *why* (17) sounds counter-intuitive. Moreover, as far as I can tell, Pollock's (DPR)-theory doesn't have any obvious way of explaining what's going on here. It sounds wrong (to my ear) to say that Pa *does* support Na , but that this support is somehow *defeated* by something else. On the contrary, it seems to me that Pa (defeasibly) *counter-supports* Na in this context. In any event, I offer this example as an invitation to clarify how Pollock's (DPR) theory works, and how it compares with (BE).

I wish I had more space to discuss other aspects of Pollock's (DPR) theory, not to mention his theory of "causal probability" and his new approach to decision theory. There is just a ton of really interesting and novel stuff in this book. And, there is also a lot of neat stuff "under the hood" that isn't (explicitly) discussed in the book (*e.g.*, some very powerful and ingenious computer programs for calculating and proving general claims about the sorts of probabilities Pollock has in mind). Working through *Thinking About Acting* was challenging and edifying. I highly recommend it to anyone interested in decision theory, probability, epistemology and/or various other related fields.

REFERENCES

- [1] Adams, E., 1975, *The Logic of Conditionals*, Dordrecht: D. Reidel.
 [2] Bennett, J., 2003, *A Philosophical Guide to Conditionals*, New York: Oxford University Press.
 [3] Carnap, R., 1962, *Logical Foundations of Probability*, 2nd edition, Chicago: University of Chicago Press.
 [4] Edgington, D., 1995, "On Conditionals", *Mind* 104(414): 235-329.
 [5] Fitelson, B., 2008, "A Decision Procedure for Probability Calculus with Applications", *Review of Symbolic Logic* 1(1): 111-125.
 [6] Garber, D., 1983, "Old Evidence and Logical Omniscience in Bayesian Confirmation Theory", in J. Earman, ed., *Testing Scientific Theories*, *Minnesota Studies in the Philosophy of Science*, volume 10, Minneapolis: University of Minnesota Press.
 [7] Hailperin, T., 1996, *Sentential Probability Logic*, Bethlehem: Lehigh University Press.
 [8] Joyce, J., 2009, "Accuracy and Coherence: Prospects for an Alethic Epistemology of Partial Belief", in Huber, F. and Schmidt-petri, C., eds., *Degrees of Belief*. New York: Springer.
 [9] Keynes, J.M., 1921, *A Treatise on Probability*, London: Macmillan.
 [10] Kolmogorov, A.N., 1956, *Foundations of Probability Theory*, New York: Chelsea.
 [11] Kolodny, N. and MacFarlane, J., 2009, "Ifs and Oughts", unpublished manuscript.
 [12] McGee, V., 1985, "A Counterexample to Modus Ponens", *Journal of Philosophy*, LXXXII (9): 462-471.
 [13] Pollock, J., 1990, *Nomic Probability and the Foundations of Induction*, New York: Oxford University Press.
 [14] ———, 2006, *Thinking About Acting*, New York: Oxford University Press.

- [15] ———, 2009, "Probable Probabilities", unpublished manuscript.
 [16] Shogenji, T., 2009, "The Degree of Epistemic Justification and the Conjunction Fallacy", to appear in *Synthese*.
 [17] Skyrms, B., 1999, *Choice and Chance: An Introduction to Inductive Logic*, 4th edition, Belmont, California: Wadsworth.
 [18] White, R., 2006, "Problems for Dogmatism", *Philosophical Studies* 131(5): 525-557.
 [19] Williamson, T., 2000, *Knowledge and its Limits*, Oxford: Oxford University Press.

PHILOSOPHY DEPARTMENT, RUTGERS UNIVERSITY, NEW BRUNSWICK, NJ 08901.

E-mail address: branden@fitelson.org

URL: <http://fitelson.org/>