

Philosophy 148 — Announcements & Such

- Overall, people did very well on the mid-term ($\mu = 90, \sigma = 16$).
- HW #2 is still being graded. We'll have those ready Thursday.
- Recall: we're using a straight grading scale for this course. Here it is:
 - A+ > 97, A (94,97], A- (90,94], B+ (87,90), B (84,87], B- (80,84], C+ (77,80], C (74,77], C- (70,74], D [50,70], F < 50.
- I've posted solutions for HW #1 (HW #2 solutions are coming soon).
- **There is a HW #3 discussion tonight @ 6pm @ 251 Dwinelle.** HW #3 is due Thursday. HW #4 will be assigned on Thursday.
- Today: a brief Review of Joyce's argument. Then, "Logical" Probability:
 - Leibnizian/Wittgensteinian Roots of "Logical" Probability
 - The Principle of Indifference
 - Carnap's (Early) Formal Theories of "Logical" Probability

Review of Joyce's Epistemic Argument for Probabilism

- Joyce aims to show that, unless q is a probability function, there will exist a q' which is closer to the truth - *no matter what the truth turns out to be*.
- The "closeness to the truth" of q in world w is based on some measure ρ of the distance between $q(X)$ and the truth-value ($w(X) = 0$ or 1) of X in w , summed over all of the X 's in S 's set of entertainable propositions \mathcal{B} .
- Joyce's argument rests on four assumptions \mathcal{A} about the inaccuracy measure I . \mathcal{A} is consistent with $\rho^\dagger(q(X), w(X)) = |q(X) - w(X)|^2$, but not $\rho^*(q(X), w(X)) = |q(X) - w(X)|$, or various other distance measures.
- Maher shows that ρ^* violates two of Joyce's axioms: (WC) and (S), and he argues that ρ^* is nonetheless a reasonable measure of "distance from truth". Maher also shows that Joyce's Theorem *fails* if one uses ρ^* .
- Joyce's defense of ρ^\dagger appeals to "Expected Distance from Truth" of q . I mentioned two problems with Joyce's reply: (1) it appeals to something that seems *pragmatic*, and (2) it *presupposes* that $q(\sim p) = 1 - q(p)$.

Whither Logical Probability? 1

- The motivation for "logical interpretations" of probability has its roots in *inductive logic*. Inductive Logic (more later) is meant to be the science of *argument strength* — a quantitative generalization of entailment (\models).
- Inductive Logic aims to explicate a quantitative measure $c(C, P)$ of the "degree to which the premises of an argument P jointly confirm its conclusion C ." Keynes, Carnap, and others worked on such theories.
- As we will study more in our confirmation unit, one of the key desiderata of IL is that c should quantitatively generalize deductive entailment:

(\mathcal{D}_1) The relations of deductive entailment and deductive refutation should be captured as limiting (extreme) values of c with cases of 'partial entailment' and 'partial refutation' lying somewhere on a c -continuum (or range) between these extreme values of the confirmation function.
- Exercises: (i) *the conditional probability* $[\Pr(C | P)]$ satisfies \mathcal{D}_1 , but *the probability of the corresponding conditional* $[\Pr(P \rightarrow C)]$ does *not*.

Whither Logical Probability? 2

- Another key historical desideratum for inductive logic is:

(\mathcal{D}_2) Inductive logic (*i.e.*, the *non*-deductive relations characterized by inductive logic) should be *objective* and *logical*.
- Carnap on this desideratum:

"Deductive logic may be regarded as the theory of the relation of logical consequence, and inductive logic as the theory of another concept [c] which is likewise objective and logical, *viz.*, ... degree of confirmation."
- Carnap on the terms "logical" and "objective" as they apply to c .

"The principal common characteristic of the statements in both fields is their independence of the contingency of facts. This characteristic justifies the application of the common term 'logic' to both fields."

"That c is objective means this: if a certain c value holds for a hypothesis (H) with respect to evidence (E), then the value $c(H, E)$ is independent of what any person may happen to think about these sentences."

Whither Logical Probability? 3

- While $\Pr(C | P)$ clearly satisfies \mathcal{D}_1 , it is unclear whether $\Pr(C | P)$ satisfies \mathcal{D}_2 . If $\mathfrak{c}(C, P) \stackrel{\text{def}}{=} \Pr(C | P)$, then, we seem to need $\Pr(C | P)$ itself to be logical if $\mathfrak{c}(C, P)$ is to be logical. Hence, the need for “logical probability”!
- Moreover, if we assume the standard definition of $\Pr(C | P)$, then we need an unconditional probability function that is itself logical. The basic “Leibnizian” idea, which underlies this “interpretation” of probability:

$$\Pr(C | P) = \frac{\Pr(P \& C)}{\Pr(P)} = \frac{\text{The proportion of logically possible worlds in which } P \& C \text{ is true}}{\text{The proportion of logically possible worlds in which } P \text{ is true}}$$
- Such unconditional logical probability functions are called logical *measure functions* (m). Intuitively, these are intended to measure the “proportion of logically possible worlds in which a proposition is true”.
- Wittgenstein, Carnap, and others give precise explications of this vague concept of “logical probability”. They work within logical *languages* \mathcal{L} , and they work with *descriptions* of possible worlds — *sentences* in \mathcal{L} .

Wittgensteinian Propositional Logical Probability 2

- Here are some important *technical* facts about Wittgensteinian \mathcal{L}_P^n -Pr:
 - Every atomic sentence in every \mathcal{L}_P^n probability model has probability $\frac{1}{2^n}$.
 - Every collection of atomic sentences $\{A_1, \dots, A_n\}$ in every \mathcal{L}_P^n probability model is mutually probabilistically independent.
 - For every pair of sentences p and q (atomic or compound) in every \mathcal{L}_P^n probability model, p and q are probabilistically dependent iff p and q are logically dependent (i.e., iff $p \models q$ or $\sim p \models q$ or $q \models p$ or $\sim q \models p$). [This depends on the *equiprobability* of the state descriptions. *Why?*]
 - For all p and q in any \mathcal{L}_P^n model, $\Pr(q | p) = 1$ if and only if $p \models q$. [This does *not* depend on the *equiprobability* of the S.D.’s. *Why?*]
- The key philosophical question about this kind of approach is:
 - What makes the assignment of *equal* probability to the *state descriptions* the “logical” probability assignment? What *logical* principle is at work here? There are 2 kinds of answers: (i) the *Principle of Indifference* (PI), and (ii) *permutation invariance*. Carnap on (PI)...

Wittgensteinian Propositional Logical Probability 1

- In the *Tractatus*, Wittgenstein presents the idea of a truth-table for a propositional language \mathcal{L}_P . He proposes a logical measure function m on \mathcal{L}_P , which assigns *equal probability* to each *state description* of \mathcal{L}_P .
- Let \mathcal{L}_P^n be a propositional language with n atomic sentences. The measure function m would assign $m(s_i) = \frac{1}{2^n}$, for all state descriptions s_i of \mathcal{L}_P^n . Here’s what m looks like over an \mathcal{L}_P^3 language (this should look familiar!):

A	B	C	State Descriptions	$m(s_i)$
T	T	T	$s_1 = A \& B \& C$	$m(s_1) = 1/8$
T	T	F	$s_2 = A \& B \& \sim C$	$m(s_2) = 1/8$
T	F	T	$s_3 = A \& \sim B \& C$	$m(s_3) = 1/8$
T	F	F	$s_4 = A \& \sim B \& \sim C$	$m(s_4) = 1/8$
F	T	T	$s_5 = \sim A \& B \& C$	$m(s_5) = 1/8$
F	T	F	$s_6 = \sim A \& B \& \sim C$	$m(s_6) = 1/8$
F	F	T	$s_7 = \sim A \& \sim B \& C$	$m(s_7) = 1/8$
F	F	F	$s_8 = \sim A \& \sim B \& \sim C$	$m(s_8) = 1/8$

Carnap on the Principle of Indifference 1

- The Principle of Indifference (PI) says, roughly, that if an epistemically rational agent’s total evidence K does not favor any member of a partition of possible states $\{s_1, \dots, s_n\}$ over any other member, then that agent’s degrees of credence (*assumed* to be probabilities!) should satisfy:

$$\Pr(s_i | K) = \Pr(s_j | K), \text{ for all } i, j.$$
- The invocation of (PI) in this context should be puzzling for a few reasons:
 - First, (PI) *sounds like* an *epistemic* principle about what an agent’s epistemic probabilities should be, under certain circumstances.
 - Should *logical* principles be determined by *epistemic* constraints? This is controversial, even for deductive logic (think inconsistent beliefs):
 - * You shouldn’t *believe everything* if your set of beliefs happens to contain a contradiction (especially, in light of the Preface!).
 - So, even for *deductive* logic, *epistemic* principles seem to make for odd *logical* constraints. Why should *inductive* logic be any different?

- Presumably, the (PI) approach goes as follows. Assume that “all we know is logic”. Call this background knowledge K_T . Assume that K_T does not favor any state description over any other. Then, m can be thought of as $\text{Pr}(\bullet | K_T)$, and we have $\text{Pr}(s_i | K_T) = \text{Pr}(s_j | K_T)$, for all i, j , as desired.
- Carnap cleverly argues that (PI) is a *logical* (not epistemic!) principle:
 - ... the statement of equiprobability to which the (PI) leads is, like all other statements of inductive probability, not a factual but a logical statement. If the knowledge of the observer does not favor any of the possible events, then with respect to this knowledge as evidence they are equiprobable.
- Here’s how the argument is supposed to go (as far as I can see):
 - Intuitively, “ K does not favor s_i over s_j (any $i \neq j$)” \Rightarrow “ K confirms s_i to the same degree as K confirms s_j (each $i \neq j$)”, i.e., $c(s_i, K) = c(s_j, K)$.
 - If we assume that $c(x, y) = \text{Pr}(x | y)$, then *logic alone* implies that:

$$c(s_i, K) = c(s_j, K) \Rightarrow \text{Pr}(s_i | K) = \text{Pr}(s_j | K), \text{ for all } i \text{ and } j.$$
 - \therefore *Logic alone* implies that if K does not favor s_i over s_j (for any $i \neq j$ in a partition of states), then $\text{Pr}(s_i | K) = \text{Pr}(s_j | K)$, for each i, j . \square

Carnapian Monadic Predicate Logical Probability 1

- Generalizing on Wittgenstein, Carnap defined his logical measure functions over sentences in monadic predicate logical languages $\mathcal{L}_Q^{m,n}$ containing n monadic predicates (F, G, \dots) and m constants (a, b, \dots).
- To fix ideas, consider the language $\mathcal{L}_Q^{2,2}$, which contains two monadic predicates F and G and two individual constants a and b .
- In $\mathcal{L}_Q^{2,2}$, we can describe 16 states, using the 16 *state descriptions* of $\mathcal{L}_Q^{2,2}$:

$Fa \ \& \ Ga \ \& \ Fb \ \& \ Gb$	$Fa \ \& \ Ga \ \& \ Fb \ \& \ \sim Gb$	$Fa \ \& \ Ga \ \& \ \sim Fb \ \& \ Gb$
$Fa \ \& \ Ga \ \& \ \sim Fb \ \& \ \sim Gb$	$Fa \ \& \ \sim Ga \ \& \ Fb \ \& \ Gb$	$Fa \ \& \ \sim Ga \ \& \ Fb \ \& \ \sim Gb$
$Fa \ \& \ \sim Ga \ \& \ \sim Fb \ \& \ Gb$	$Fa \ \& \ \sim Ga \ \& \ \sim Fb \ \& \ \sim Gb$	$\sim Fa \ \& \ Ga \ \& \ Fb \ \& \ Gb$
$\sim Fa \ \& \ Ga \ \& \ Fb \ \& \ \sim Gb$	$\sim Fa \ \& \ Ga \ \& \ \sim Fb \ \& \ Gb$	$\sim Fa \ \& \ Ga \ \& \ \sim Fb \ \& \ \sim Gb$
$\sim Fa \ \& \ \sim Ga \ \& \ Fb \ \& \ Gb$	$\sim Fa \ \& \ \sim Ga \ \& \ Fb \ \& \ \sim Gb$	$\sim Fa \ \& \ \sim Ga \ \& \ \sim Fb \ \& \ Gb$
$\sim Fa \ \& \ \sim Ga \ \& \ \sim Fb \ \& \ \sim Gb$		

Carnap on the Principle of Indifference 2

- Carnap’s argument is clever, but ultimately not terribly compelling.
- As we will see next, Carnap’s own Predicate-Logical approach to “logical probability” deviates from the application of (PI) to the state descriptions of his logical languages [and, for reasons that look *epistemic*, yet again!].
- Aside from these worries about the “logicality” of (PI) and its success at generating the “right ‘logical’ probability models”, there are other problems with Carnap’s argument. Here’s a cursory sketch (more later):
 - Carnap *assumes* that the correct (probabilistic) explication of the *confirmation* function is just the conditional probability function, i.e.,

$$c(x, y) = \text{Pr}(x | y)$$
 - If it were obvious that $\text{Pr}(x | y)$ is the best (probabilistic) explication of $c(x, y)$, then Carnap’s argument might be sound (if not informative!).
 - We’ll soon see that $\text{Pr}(x | y)$ may *not* be the best explication of $c(x, y)$.
- We’ll discuss other aspects of the (PI) in the next units of the course.

Carnapian Monadic Predicate Logical Probability 2

- Following (PI), Carnap’s first measure function m^\dagger assigns *equal probability* to each state description s_i of $\mathcal{L}_Q^{m,n}$. In our $\mathcal{L}_Q^{2,2}$, $m^\dagger(s_i) = \frac{1}{16}$.
- We extend m^\dagger to all $p \in \mathcal{L}_Q^{m,n}$ in the standard way (the Pr of a disjunction of mutually exclusive sentences is the sum of the Pr’s of its disjuncts).
- Since every $p \in \mathcal{L}_Q^{m,n}$ is equivalent to some disjunction of state descriptions, and every pair of state descriptions is mutually exclusive, this gives us a complete unconditional Pr-function $\text{Pr}^\dagger(\cdot)$ over $\mathcal{L}_Q^{m,n}$.
- Finally, we define the conditional probability function $\text{Pr}^\dagger(q | p)$ over pairs of sentences in $\mathcal{L}_Q^{m,n}$ (in the standard way) as the following *ratio*: $\frac{\text{Pr}^\dagger(p \ \& \ q)}{\text{Pr}^\dagger(p)}$.
- Claims of the form ‘ $\text{Pr}^\dagger(q | p) = x$ ’ are *analytic in $\mathcal{L}_Q^{m,n}$* since their truth-values are determined solely by the syntactical structure of $\mathcal{L}_Q^{m,n}$.
- But, why is *this* choice of measure function m^\dagger *logical*? Logicality is ensured by the application of (PI) to state descriptions. Or is it?

Carnapian Monadic Predicate Logical Probability 3

- As it turns out, Carnap ultimately *rejects* the measure function m^\dagger in favor of an alternative measure function m^* , for epistemic-sounding reasons.
- Carnap notes that m^\dagger causes Pr^\dagger to have the following property ($b \neq a$):

$$(*) \quad Pr^\dagger(Fb | Fa) = \frac{Pr^\dagger(Fb \& Fa)}{Pr^\dagger(Fa)} = \frac{4 \cdot \frac{1}{16}}{8 \cdot \frac{1}{16}} = \frac{1}{2} = 8 \cdot \frac{1}{16} = Pr^\dagger(Fb)$$

- In other words, (*) says that one object a 's having property F can never raise the probability that another object b also has F . And, this generalizes to any number of F s: $Pr^\dagger(Fa_1 | Fa_2 \& \dots \& Fa_m) = Pr^\dagger(Fa_1)$.
- Carnap (*et al*) characterize this (in *epistemic* terms) as m^\dagger leading to a function Pr^\dagger that fails to allow for "learning from experience". This is *epistemic and diachronic*, since it assumes *learning-by-conditionalization*.
- Carnap views this consequence of applying (PI) to the state descriptions of $\mathcal{L}_Q^{m,n}$ as unacceptable. Does he then think that K_T "favors" some state descriptions over others. But, which ones? Enter the m^* measure ...

Carnapian Monadic Predicate Logical Probability 5

- We can then define $Pr^*(\bullet)$ and $Pr^*(\bullet | \bullet)$ in terms of m^* , by assuming equiprobability of *states within* structure descriptions. Let's compare the m^\dagger and m^* distributions over the 16 state descriptions of $\mathcal{L}_Q^{2,2}$:

Fa	Ga	Fb	Gb	State Descriptions (s_i)	$m^\dagger(s_i)$	$m^*(s_i)$
T	T	T	T	$Fa \& Ga \& Fb \& Gb$	1/16	1/10
T	T	T	F	$Fa \& Ga \& Fb \& \sim Gb$	1/16	1/20
T	T	F	T	$Fa \& Ga \& \sim Fb \& Gb$	1/16	1/20
T	T	F	F	$Fa \& Ga \& \sim Fb \& \sim Gb$	1/16	1/20
T	F	T	T	$Fa \& \sim Ga \& Fb \& Gb$	1/16	1/20
T	F	T	F	$Fa \& \sim Ga \& Fb \& \sim Gb$	1/16	1/10
T	F	F	T	$Fa \& \sim Ga \& \sim Fb \& Gb$	1/16	1/20
T	F	F	F	$Fa \& \sim Ga \& \sim Fb \& \sim Gb$	1/16	1/20
F	T	T	T	$\sim Fa \& Ga \& Fb \& Gb$	1/16	1/20
F	T	T	F	$\sim Fa \& Ga \& Fb \& \sim Gb$	1/16	1/20
F	T	F	T	$\sim Fa \& Ga \& \sim Fb \& Gb$	1/16	1/10
F	T	F	F	$\sim Fa \& Ga \& \sim Fb \& \sim Gb$	1/16	1/20
F	F	T	T	$\sim Fa \& \sim Ga \& Fb \& Gb$	1/16	1/20
F	F	T	F	$\sim Fa \& \sim Ga \& Fb \& \sim Gb$	1/16	1/20
F	F	F	T	$\sim Fa \& \sim Ga \& \sim Fb \& Gb$	1/16	1/20
F	F	F	F	$\sim Fa \& \sim Ga \& \sim Fb \& \sim Gb$	1/16	1/10

- Now, m^* does *not* have the "no learning from experience" property (*):

Carnapian Monadic Predicate Logical Probability 4

- Two state descriptions s_i and s_j in $\mathcal{L}_Q^{m,n}$ are *permutations* of each other if one can be obtained from the other by a permutation of constants.
- " $Fa \& \sim Ga \& \sim Fb \& Gb$ " can be obtained from " $\sim Fa \& Ga \& Fb \& \sim Gb$ " by permuting " a " and " b ". Thus, " $Fa \& \sim Ga \& \sim Fb \& Gb$ " and " $\sim Fa \& Ga \& Fb \& \sim Gb$ " are permutations of each other (in $\mathcal{L}_Q^{2,2}$).
- A *structure description* is a disjunction of state descriptions, each of which is a permutation of the others. $\mathcal{L}_Q^{2,2}$ has 10 structure descriptions:

$(Fa \& \sim Ga \& \sim Fb \& Gb) \vee (\sim Fa \& Ga \& Fb \& \sim Gb)$	$Fa \& \sim Ga \& Fb \& \sim Gb$
$(Fa \& Ga \& Fb \& \sim Gb) \vee (Fa \& \sim Ga \& Fb \& Gb)$	$Fa \& Ga \& Fb \& Gb$
$(Fa \& Ga \& \sim Fb \& Gb) \vee (\sim Fa \& Ga \& Fb \& Gb)$	$\sim Fa \& Ga \& \sim Fb \& Gb$
$(Fa \& Ga \& \sim Fb \& \sim Gb) \vee (\sim Fa \& \sim Ga \& Fb \& Gb)$	$\sim Fa \& \sim Ga \& \sim Fb \& \sim Gb$
$(Fa \& \sim Ga \& \sim Fb \& \sim Gb) \vee (\sim Fa \& \sim Ga \& Fb \& \sim Gb)$	
$(\sim Fa \& Ga \& \sim Fb \& \sim Gb) \vee (\sim Fa \& \sim Ga \& \sim Fb \& Gb)$	
- The measure m^* [\cdot : Pr^*] assigns equal probability to *structure descriptions*. m^* takes individuals to be *indistinguishable* (analogy: Bose-Einstein statistics in QM), and m^\dagger does not (analogy: Fermi-Dirac statistics in QM). Otherwise, they operate in the same way.

$$Pr^*(Fb | Fa) = \frac{Pr^*(Fb \& Fa)}{Pr^*(Fa)} = \frac{2 \cdot \frac{1}{10} + 2 \cdot \frac{1}{20}}{2 \cdot \frac{1}{10} + 6 \cdot \frac{1}{20}} = \frac{3}{5} > 2 \cdot \frac{1}{10} + 6 \cdot \frac{1}{20} = \frac{1}{2} = Pr^*(Fb)$$

- Generally, $Pr^*(\bullet | \bullet)$ says that the more objects that are assumed to have F , the more probable it is that other objects will also have F . This is called *instantial relevance*. Carnap prefers Pr^* over Pr^\dagger for this reason.
- Is Carnap committed to the view that K_T favors certain state descriptions over others? He prefers $m^*(Fa \& Ga \& Fb \& Gb) > m^*(Fa \& Ga \& Fb \& \sim Gb)$ to $m^\dagger(Fa \& Ga \& Fb \& Gb) = m^\dagger(Fa \& Ga \& Fb \& \sim Gb)$. But, *why*?
- Carnap realizes that if a theory of "logical probability" is to provide a foundation for claims about *evidence*, then it *must* be able to furnish Pr -models with *correlations* between *logically independent* claims.
- This is the tip of the iceberg for linguistic theories of "logical probability" like Wittgenstein's and Carnap's. Such theories are unable to emulate a broad enough range of probability functions, so as (D_3) to be generally *applicable* for modeling evidential relations in epistemic contexts.
- We will return to this *applicability* issue later in the course...

Carnapian Logical Probability: Analogy and Similarity 1

- Moreover, Pr^* has other shortcomings. Carnap and others have endorsed the following “*analogical*” principle: *a* and *b* differing regarding *F* should *weaken* the inference from *Ga* to *Gb*, *without completely undermining it*:

$$Pr(Gb | Ga) > Pr(Gb | Fa \& Ga \& \sim Fb) > Pr(Gb)$$

Neither Pr^* nor Pr^\dagger satisfies this condition. This will be a HW #3 problem.


- Generally: if *a* has properties $P_1 \dots P_n$, and *b* has $P_1 \dots P_{n-2}$, but lacks P_{n-1} , that should be of *some* relevance to *b*’s having P_n . In the $n = 3$ case:
 $Pr(Hb | Ha) > Pr(Hb | Fa \& Ga \& Ha \& Fb \& \sim Gb)$

$$> Pr(Hb | Fa \& Ga \& Ha \& \sim Fb \& \sim Gb) > Pr(Hb)$$

- I.e.*, Differing on 2 properties should be worse than 1, but neither should completely undermine instantial relevance. Pr^\dagger and Pr^* violate this.
- Note: this measures “similarity” by “shared predicate counting”. This is unfortunate, as it leads to Pr -functions that are *language-relative*...

Carnapian Logical Probability: Analogy and Similarity 3

- That is, “the degree of similarity of *a* and *b*” depends sensitively on the *syntax* of the language one uses to *describe a* and *b*. Here’s why.
- The *ABCD* language consists of four predicates *A*, *B*, *C*, and *D*. And, the *XYZU* language also has four predicates *X*, *Y*, *Z*, and *U* such that $Xx \models Ax \equiv Bx$, $Yx \models Bx \equiv Cx$, $Zx \models Cx \equiv Ax$, and $Ux \models Dx$.

 *ABCD* and *XYZU* are (extra-systematically) **expressively equivalent**.

Anything that can be said in *ABCD* can be said in *XYZU*, and conversely — *intuitively, there is no semantic difference between the two languages*.

- Now, consider two objects *a* and *b* such that:

$$Aa \& Ba \& Ca \& Da$$

$$Ab \& \sim Bb \& Cb \& Db$$

- Question: How similar are *a* and *b* in our “predicate-sharing” sense?

Carnapian Logical Probability: Analogy and Similarity 2

- Let x_1, \dots, x_n be the objects that fall under the predicate *X* [*i.e.*, those in $Ext(X)$]. And, let $s(x_1, \dots, x_n)$ be some measure of “the degree to which the objects falling under *X* are similar to each other”.
- Carnap doesn’t offer much in the way of a *theory* of $s(x_1, \dots, x_n)$. But, his discussion suggests the following account of $s(x_1, \dots, x_n)$:
- Let $\mathcal{P}(x)$ be the set of predicates that *x* falls under. Then, define:

$$s(x_1, \dots, x_n) = \left| \bigcap_i \mathcal{P}(x_i) \right|$$

- That is, $s(x_1, \dots, x_n)$ is the size (cardinality) of the intersection of all the $\mathcal{P}(x_i)$. This is “the size of the set of shared predicates of the x_i ”.
- There is a problem with this idea. Next, I will present an argument which shows that this measure of similarity is *language variant*.

Carnapian Logical Probability: Analogy and Similarity 4

- Answer: That depends on which of our expressively equivalent languages we use to describe *a* and *b*! To see this, note that in *XYZU* we have:

$$Xa \& Ya \& Za \& Ua$$

$$\sim Xb \& \sim Yb \& Zb \& Ub$$

- Therefore, in *ABCD*, *a* and *b* share three predicates. But, in *XYZU*, *a* and *b* share only two predicates. Or, to use a modified notation, we have:

$$s_{ABCD}(a, b) = 3 \neq 2 = s_{XYZU}(a, b)$$

- Not good. Probabilities should *not* be language-variant. It shouldn’t matter which language you use to describe the world — semantically equivalent statements should be probabilistically indistinguishable.
- One consequence is that if $p \models q$, $x \models y$ and $z \models u$, then we shouldn’t have both $Pr(p | x) > Pr(p | z)$ and $Pr(q | u) > Pr(q | y)$. Carnapian principles of analogy and similarity *contradict* this requirement.

Carnapian Logical Probability: Analogy and Similarity 5

- Here's a concise way of stating the general Carnapian analogical principle:

(A) If $n > m$, then $\Pr(Xa \mid Xb \ \& \ s(a, b) = n) > \Pr(Xa \mid Xb \ \& \ s(a, b) = m)$

- Applying this principle to our example yields *both* of the following:

(1) $\Pr(Da \mid Aa \ \& \ Ba \ \& \ Ca \ \& \ Ab \ \& \ \sim Bb \ \& \ Cb) > \Pr(Da \mid Aa \ \& \ Ba \ \& \ Ca \ \& \ Ab \ \& \ \sim Bb \ \& \ \sim Cb)$

(2) $\Pr(Ua \mid Xa \ \& \ Ya \ \& \ Za \ \& \ \sim Xb \ \& \ Yb \ \& \ Zb) > \Pr(Ua \mid Xa \ \& \ Ya \ \& \ Za \ \& \ \sim Xb \ \& \ \sim Yb \ \& \ Zb)$

- Now, let $p \stackrel{\text{def}}{=} Da$, $q \stackrel{\text{def}}{=} Ua$, and

$x \stackrel{\text{def}}{=} Aa \ \& \ Ba \ \& \ Ca \ \& \ Ab \ \& \ \sim Bb \ \& \ Cb$, $y \stackrel{\text{def}}{=} Xa \ \& \ Ya \ \& \ Za \ \& \ \sim Xb \ \& \ \sim Yb \ \& \ Zb$,

$z \stackrel{\text{def}}{=} Xa \ \& \ Ya \ \& \ Za \ \& \ \sim Xb \ \& \ Yb \ \& \ Zb$, $u \stackrel{\text{def}}{=} Aa \ \& \ Ba \ \& \ Ca \ \& \ Ab \ \& \ \sim Bb \ \& \ \sim Cb$.

- Then, $p \models q$, $x \models y$ and $z \models u$, but the Carnapian principle (A) implies *both* (1) $\Pr(p \mid x) > \Pr(p \mid u)$, *and* (2) $\Pr(q \mid z) > \Pr(q \mid y)$. Bad.
- It seems that principle (A) must go. Otherwise, some restriction on the choice of language is required to block inferring both (1) and (2) from it.