

Philosophy 148 — In-Class Quiz Answer Key

02/14/08

(1) Write the three (Kolmogorov) probability axioms we are using in this class:

**A:** If  $\Pr(\bullet)$  is defined over a language  $\mathcal{L}$ , then for all  $p, q \in \mathcal{L}$ :

1.  $\Pr(p) \geq 0$ .
2. If  $p \models \top$ , then  $\Pr(p) = 1$ .
3. If  $p \& q \models \perp$ , then  $\Pr(p \vee q) = \Pr(p) + \Pr(q)$ .

(2) Write our definition of the conditional probability  $\Pr(X | Y)$ :

**A:**  $\Pr(X | Y) \stackrel{\text{def}}{=} \frac{\Pr(X \& Y)}{\Pr(Y)}$

(3)  $\sim(X \rightarrow Y) \models X \& \sim Y$

T/ F

**A:** T.  $X \rightarrow Y$  is true on every interpretation but the one in which  $X$  is true and  $Y$  is false. So  $\sim(X \rightarrow Y)$  is true just in case  $X \& \sim Y$ .

(4) Consider these two statements:  $p \equiv q$  and  $p \& \sim q$ .

(a) These two statements are inconsistent (mutually exclusive).

T/ F

(b) These two statements are contradictory.

T/ F

**A:** T, F. Consider the truth-table below. There is no line on which both statements are true; therefore, they are inconsistent. However, they do not always have opposite truth-values — see line 3. So they are not contradictory.

$p$	$q$	$p \equiv q$	$p \& \sim q$
T	T	T	F
T	F	F	T
F	T	F	F
F	F	T	F

(5) Consider a monadic predicate-logical language  $\mathcal{L}$  with two constants  $a$  and  $b$  (think: a universe of discourse containing two objects) and two predicates  $F$  and  $G$ . Which of the following state descriptions of  $\mathcal{L}$  is entailed by the universal claim  $(\forall x)(Fx \& \sim Gx)$ ? Circle the correct answer. (Exactly one is correct.)

(i)  $Fa \& Ga \& Fb \& Gb$

(iii)  $Fa \& \sim Ga \& Fb \& \sim Gb$

(ii)  $Fa \& Ga \& \sim Fb \& \sim Gb$

(iv)  $\sim Fa \& \sim Ga \& \sim Fb \& \sim Gb$

**A:** (iii). The universal quantification says that for any object in the universe of discourse,  $F$  is true of that object and  $G$  is false of it.

(6) Consider the probability model  $\mathcal{M}$  described in this stochastic truth-table:

$X$	$Y$	State	$\Pr(s_i)$
T	T	$s_1$	0.1
T	F	$s_2$	0.2
F	T	$s_3$	0.3
F	F	$s_4$	0.4

Solve the following problems, concerning this model:

(a) Calculate the value of  $\Pr(X)$  in  $\mathcal{M}$ .

**A:** 0.3.  $\Pr(X) = \Pr(X \& Y) + \Pr(X \& \sim Y) = 0.1 + 0.2 = 0.3$

(b) Calculate the value of  $\Pr(X | Y)$  in  $\mathcal{M}$ . **A:**  $\frac{1}{4}$ .

$$\Pr(X | Y) = \frac{\Pr(X \& Y)}{\Pr(Y)} = \frac{\Pr(X \& Y)}{\Pr(X \& Y) + \Pr(\sim X \& Y)} = \frac{0.1}{0.1 + 0.3} = \frac{1}{4}$$


(c)  $\mathcal{M}$  provides a counter-example to  $\Pr[\sim(X \& Y)] > \Pr(\sim X \rightarrow Y)$ . T/ F

**A:** F.  $\Pr(X \& Y) = 0.1$ , so  $\Pr[\sim(X \& Y)] = 0.9$ .

$\sim X \rightarrow Y$  is false just in case  $\sim X \& \sim Y$ .  $\Pr(\sim X \& \sim Y) = 0.4$ , so  $\Pr(\sim X \rightarrow Y) = 0.6$ .

Thus *for this model*  $\Pr[\sim(X \& Y)] > \Pr(\sim X \rightarrow Y)$ , so *this model* does *not* provide a counter-example to the inequality. Note that this does not show the inequality is true for *every* model.

(7) Consider the probability model  $\mathcal{M}'$  depicted by the following Stochastic Venn Diagram (note: the diagram IS drawn to scale, with areas of regions proportional to probabilities of corresponding propositions in  $\mathcal{M}'$ ):



Circle true or false for each of the following, as they pertain to  $\mathcal{M}'$ :

(a)  $X \vDash Y$  **A:** F. There are points in the  $X$  region where  $Y$  is not true.

(b)  $Y \vDash X$  **A:** T. At every point in the  $Y$  region,  $X$  is true.

(c)  $\Pr(X | Y) > 0.5$  **A:** T. Because  $Y \vDash X$ ,  $\Pr(X | Y) = 1$ .

(d)  $\Pr(Y | X) > 0.5$  **A:** F. The  $Y$  region occupies less than half of the  $X$  region.

(e)  $X$  and  $Y$  are correlated. **A:** T. As we just saw,  $\Pr(X | Y) = 1$ . But  $\Pr(X) < 1$  (because  $X$  does not occupy the entire rectangle). So  $\Pr(X | Y) > \Pr(X)$ ;  $X$  and  $Y$  are correlated.

(8) Suppose I roll a fair six-sided die (equal chance of any face coming up) and then flip a fair coin (equal chance of each side coming up), with the outcome of the die roll independent of the outcome of the coin flip. Define statements  $A$ ,  $B$ , and  $C$  as follows:

$A =$  'The coin came up heads'  
 $B =$  'The die roll came up 3'  
 $C =$  'The die roll came up with an odd number'

Assuming  $\Pr$  is a probability function in some probability model compatible with the above description of the situation, answer the following questions:

(a) What is  $\Pr(A \& B)$ ? **A:**  $\frac{1}{12}$ .

$A$  and  $B$  are independent, so  $\Pr(A \& B) = \Pr(A) \cdot \Pr(B) = \frac{1}{2} \cdot \frac{1}{6} = \frac{1}{12}$ .

(b) What is  $\Pr(A \vee C)$ ? **A:**  $\frac{3}{4}$ . By the general additivity theorem,  $\Pr(A \vee C) = \Pr(A) + \Pr(C) - \Pr(A \& C)$ .  
By independence,  $\Pr(A \& C) = \Pr(A) \cdot \Pr(C)$ . So

$$\Pr(A \vee C) = \Pr(A) + \Pr(C) - \Pr(A) \cdot \Pr(C) = \frac{1}{2} + \frac{1}{2} - \frac{1}{2} \cdot \frac{1}{2} = \frac{3}{4}$$

(c) What is  $\Pr(B | C)$ ? **A:**  $\frac{1}{3}$ .  $\Pr(B | C) = \frac{\Pr(B \& C)}{\Pr(C)} = \frac{1/6}{1/2} = \frac{1}{3}$

(d) What is  $\Pr(C | A)$ ? **A:**  $\frac{1}{2}$ . By independence,  $\Pr(C | A) = \Pr(C) = \frac{1}{2}$ .